

Lower Bounds for  $K_2^{\text{top}}(\hat{\mathbb{Z}}_p\pi)$  and  $K_2(\mathbb{Z}\pi)$ 

ROBERT OLIVER

*Matematisk Institut, Ny Munkegade, DK-8000 Aarhus C, Denmark**Communicated by A. Fröhlich*

Received February 1, 1983

As seen in [11] and [4], the groups  $K_2(\hat{\mathbb{Z}}_p\pi)$  ( $p$  a prime and  $\pi$  a finite group) play a role both in understanding  $SK_1(\mathbb{Z}\pi)$  and in getting lower estimates for the size of  $K_2(\mathbb{Z}\pi)$  or  $Wh_2(\mathbb{Z}\pi)$ . In particular, it was shown in [11] that the leading problem for computing  $SK_1(\mathbb{Z}\pi)$  for arbitrary finite  $\pi$  is to understand  $K_2(\hat{\mathbb{Z}}_p\pi)$  for all primes  $p$  and  $p$ -groups  $\pi$ . In this paper, we detect elements in  $K_2(\hat{\mathbb{Z}}_p\pi)$  by describing the groups

$$K_2^*(\hat{\mathbb{Z}}_p\pi) = \varprojlim \text{Coker}[K_2(\hat{\mathbb{Z}}_p\tilde{\pi}) \rightarrow K_2(\hat{\mathbb{Z}}_p\pi)]$$

and

$$Wh_2^*(\hat{\mathbb{Z}}_p\pi) = \varprojlim \text{Coker}[Wh_2(\hat{\mathbb{Z}}_p\tilde{\pi}) \rightarrow Wh_2(\hat{\mathbb{Z}}_p\pi)],$$

where the limits are taken over all surjections of  $p$ -groups  $\tilde{\pi} \twoheadrightarrow \pi$ . In other words, we look at  $K_2(\hat{\mathbb{Z}}_p\pi)$  and  $Wh_2(\hat{\mathbb{Z}}_p\pi)$  after dividing out by all elements coming from arbitrarily large  $p$ -groups mapping onto  $\pi$ .

More generally, we let  $A$  be any unramified  $p$ -ring—the ring of integers in any finite unramified extension  $K$  of  $\hat{\mathbb{Q}}_p$ —and study the groups  $K_2^*(A\pi)$  and  $Wh_2^*(A\pi)$  defined analogously for any  $p$ -group  $\pi$ . In order to state the main result, we define

$$\mathcal{U}(A\pi) = H_1(\pi; A\pi) / \langle g \otimes \lambda g^n : g \in \pi, \lambda \in A, n \in \mathbb{Z} \rangle,$$

where  $\pi$  acts on  $A\pi$  via conjugation. Then

$$\mathcal{U}(A\pi) \cong \sum_{i=1}^k (Z_\pi(g_i)/S_\pi(g_i))^{ab} \otimes A(g_i),$$

where  $g_1, \dots, g_k$  are conjugacy class representatives for  $\pi$ ,  $Z_\pi(g_i)$  is the centralizer, and

$$S_\pi(g_i) = \langle h \in \pi : h^n = g_i, \text{ some } n \rangle.$$

There is a map

$$v: \mathcal{U}(A\pi) \rightarrow H_2(\pi)$$

defined by setting  $v(h \otimes \lambda g) = \text{Tr}(\lambda) \cdot (h \wedge g)$  (where  $h \wedge g \in H_2(\pi)$  is an element defined for each pair of commuting elements  $h, g \in \pi$ , and  $\text{Tr}: A \rightarrow \hat{\mathbb{Z}}_p$  is the trace map). Finally, we set

$$\mathcal{U}_0(A\pi) = \text{Ker}[v: \mathcal{U}(A\pi) \rightarrow H_2(\pi)].$$

The main result is the following (Theorems 3.6 and 4.2 and Proposition 3.8):

**THEOREM.** *For any prime  $p$ , any unramified  $p$ -ring  $A$ , and any  $p$ -group  $\pi$ , there is a natural exact sequence*

$$H_3(\pi) \xrightarrow{\eta_{A\pi}} Wh_2^*(A\pi) \xrightarrow{\Gamma_2^*} \mathcal{U}_0(A\pi) \rightarrow 0, \quad (1)$$

with the properties:

- (i)  $\Gamma_2(\{g, u\}) = g \otimes \Gamma(u)$  for all  $g \in \pi$  and  $u \in 1 + I(\hat{\mathbb{Z}}_p[Z_\pi(g)])$  (where  $\Gamma$  is the map defined in [10] and  $I$  denotes augmentation ideal).
- (ii)  $\text{Ker}(\eta_{A\pi})$  is independent of  $A$ .
- (iii)  $\eta_{A\pi} = 0$  if  $\pi$  is abelian (and so  $Wh_2^*(A\pi) \cong \mathcal{U}_0(A\pi)$ ).

Furthermore, there is a short exact sequence

$$0 \rightarrow H_2(\pi) \rightarrow K_2^*(A\pi) \rightarrow Wh_2^*(A\pi) \rightarrow 0$$

which is naturally split when  $A = \hat{\mathbb{Z}}_p$ .

The most important point here is the existence of the surjection  $\Gamma_2^*$  having property (i). This is the basis for most of the applications in Sections 5, 6, and 7.

The significance of the contribution of  $H_3(\pi)$  to  $Wh_2(A\pi)$  is still a mystery. A precise but complicated description of  $\text{Ker}(\eta_{A\pi})$  (and also of the extension) is given in Theorem 3.6. This can in particular be used to construct examples for any  $p$  of  $p$ -groups  $\pi$  where  $\eta_{A\pi} \neq 0$  (Proposition 4.3). But we have been unable to find any simple way of describing  $\text{Im}(\eta_{A\pi})$ .

In particular, these results show that among  $p$ -groups  $\pi$ ,  $K_2^*(\hat{\mathbb{Z}}_p\pi) = 0$  if and only if  $\pi$  is cyclic or quaternionic. That any surjection of a  $p$ -group onto  $\mathbb{Z}/p^n$  induces a surjection onto  $K_2(\hat{\mathbb{Z}}_p[\mathbb{Z}/p^n])$  is well known, but that the corresponding result for quaternionic groups holds seems surprising.

Another interesting consequence comes from combining these results with those in [10]. The image of

$$v: \mathcal{U}(A\pi) \rightarrow H_2(\pi)$$

is the subgroup generated by  $h \wedge g$  for all commuting  $h, g \in \pi$ , and this was seen in [10] to be equal to  $H_2^{ab}(\pi)$ . We can also identify the group  $\overline{I}(A\pi)$

(see [10]:  $I(A\pi)$  denotes the augmentation ideal) with  $H_0(\pi; I(A\pi))$ . Sequence (1) above now combines with Theorems 2 and 3 in [10] to give:

**THEOREM.** *For any unramified  $p$ -ring  $A$  and any  $p$ -group  $\pi$ , there is an exact sequence*

$$\begin{aligned} H_3(\pi) &\xrightarrow{\eta_{A\pi}} Wh_2^*(A\pi) \xrightarrow{I_2^*} H_1(\pi; I(A\pi)) / \langle g \otimes \lambda(g^n - 1) \rangle \rightarrow H_2(\pi) \\ &\longrightarrow Wh(A\pi) \xrightarrow{I} H_0(\pi; I(A\pi)) \rightarrow H_1(\pi) \rightarrow 0. \end{aligned}$$

An obvious question is whether this can be lifted to some exact sequence involving  $Wh_2^{\text{top}}(A\pi)$ .

We start in Section 0 with a brief sketch of the construction of sequence (1) above. This is intended to help clarify and make less mysterious the details given later in Sections 2 and 3. Also, at the end of Section 0, we show that the projection  $K_2(A\pi) \rightarrow K_2^*(A\pi)$  factors through

$$K_2^{\text{top}}(A\pi) = \varinjlim_n K_2(A/p^n[\pi]),$$

i.e., that  $K_2^*(A\pi)$  can be regarded as a quotient of  $K_2^{\text{top}}(A\pi)$ .

Section 1 deals with the main technical lemma behind the computation, and this is used in Section 2 to describe  $K_1(A\tilde{\pi}, I)$  when  $\tilde{\pi} \twoheadrightarrow \pi$  is a surjection of  $p$ -groups and  $I = \text{Ker}[A\tilde{\pi} \rightarrow A\pi]$ . The formulas for  $K_2^*(A\pi)$  and  $Wh_2^*(A\pi)$  are derived in Section 3, and the map  $\eta_{A\pi}$  in the above theorems is studied in Section 4.

The last three sections give some applications of these results. In Section 5, the surjectivity of  $I_2^*$  is combined with localization sequences to get lower bounds on the size of  $K_2(\mathbb{Z}\pi)$  and  $Wh_2(\pi)$ . This extends results of Dennis, Keating, and Stein in [4], results which helped to motivate much of this work.

In Section 6, the analogous groups  $K_2^*(A/p^n[\pi])$  are defined, and described in terms of  $H_2(\pi)$  and  $\mathcal{U}_0(A\pi)$ . The main result there is that

$$K_2(A/p[\pi]) = K_2^*(A/p[\pi])$$

for an abelian  $p$ -group  $\pi$ : this yields an explicit description of  $K_2(F\pi)$  when  $F$  is any finite field and  $\pi$  any finite abelian group. This result also gives encouragement that  $K_2^*(A\pi)$  detects a significant portion of  $K_2^{\text{top}}(A\pi)$ . The hope, of course, is that the kernel

$$\text{Ker}[K_2^{\text{top}}(A\pi) \rightarrow K_2^*(A\pi)]$$

will turn out to be generated by an easily describable set of elements. The best possible result would be that it is generated by symbols  $\{r, x\}$  for  $r \in A^*$  and  $x \in (A\pi)^*$ , but this is probably too much to hope for.

Finally, in Section 7, applications are given toward the computation of  $SK_1(\mathbb{Z}\pi)$  or  $Cl_1(\mathbb{Z}\pi)$  for  $p$ -groups  $\pi$ . The results on  $K_2^*(\hat{\mathbb{Z}}_p\pi)$  are used to get upper and lower bounds for  $Cl_1(\mathbb{Z}\pi)$ , as well as to motivate a conjecture for a combinatorial algorithm for computing  $Cl_1(\mathbb{Z}\pi)$  when  $\pi$  is a  $p$ -group and  $p$  is odd. As an example,  $SK_1(\mathbb{Z}\pi)$  is then computed when  $\pi$  is a non-abelian group of order  $p^3$  (and  $p$  any odd prime).

## 0

We start by sketching a quick construction of the exact sequence

$$H_3(\pi) \rightarrow Wh_2^*(A\pi) \rightarrow \mathcal{U}(A\pi) \xrightarrow{v} H_2(\pi).$$

This is much simpler than the construction to be given in detail in Sections 2 and 3, but it gives no information on the size of the image of  $H_3(\pi)$  in  $Wh_2(A\pi)$ , nor on how the extension at  $Wh_2^*(A\pi)$  behaves.

Throughout this section, we let  $A$  be a fixed unramified  $p$ -ring (i.e., the ring of integers in some finite unramified extension of  $\hat{\mathbb{Q}}_p$ ), and let  $\pi$  be a fixed  $p$ -group. For any surjection  $\alpha: \tilde{\pi} \twoheadrightarrow \pi$  of  $p$ -groups, define a group  $G$  and an order  $\mathfrak{U}$  as pullbacks

$$\begin{array}{ccc} G & \xrightarrow{r_1} & \tilde{\pi} \\ \downarrow r_2 & (1) & \downarrow \alpha \\ \tilde{\pi} & \xrightarrow{\alpha} & \pi \end{array} \quad \text{and} \quad \begin{array}{ccc} \mathfrak{U} & \xrightarrow{r_1} & A\tilde{\pi} \\ \downarrow r_2 & (2) & \downarrow \\ A\tilde{\pi} & \longrightarrow & A\pi \end{array}$$

We regard  $G$ ,  $\mathfrak{U}$ , and  $\tilde{\pi}$  as depending on  $\alpha$ . Set

$$I_i = \text{Ker}[r_i: AG \rightarrow A\tilde{\pi}] \quad (i = 1, 2)$$

and let  $\psi: AG \twoheadrightarrow \mathfrak{U}$  be the obvious surjection. Then  $\text{Ker}(\psi) = I_1 \cap I_2 = I_1 I_2$  (see Lemma 2.4 below); so  $\mathfrak{U} \cong AG/I_1 I_2$ , and

$$K_1(\mathfrak{U}) \cong K_1(AG)/(1 + I_1 I_2). \quad (3)$$

The five-term homology sequences for the extensions

$$1 \rightarrow \rho_1 \rightarrow G \xrightarrow{r_1} \tilde{\pi} \rightarrow 1 \quad \text{and} \quad 1 \rightarrow \rho \rightarrow \tilde{\pi} \xrightarrow{\alpha} \pi \rightarrow 1$$

(see Proposition 2.1) induce a Mayer-Vietoris exact sequence

$$H_2(\tilde{\pi}) \oplus H_2(\tilde{\pi}) \rightarrow H_2(\pi) \rightarrow G^{ab} \rightarrow \tilde{\pi}^{ab} \oplus \tilde{\pi}^{ab} \rightarrow \pi^{ab} \rightarrow 1 \quad (4)$$

(note that  $\rho_1 \cong \rho$  and  $\rho_1/[G, \rho_1] \cong \rho/[\tilde{\pi}, \rho]$ ). Upon comparing this with

the  $K$ -theory exact sequence for (2), we get an exact sequence of Whitehead groups:

$$\begin{aligned} Wh_2(A\tilde{\pi}) \oplus Wh_2(A\tilde{\pi}) &\longrightarrow Wh_2(A\pi) \xrightarrow{\partial} \frac{Wh(AG)}{1 + I_1 I_2} \\ &\xrightarrow{r_1 * \oplus r_2 *} Wh(A\tilde{\pi}) \oplus Wh(A\tilde{\pi}). \end{aligned} \quad (5)$$

In other words,

$$\begin{aligned} \text{Coker}[\alpha_*: Wh_2(A\tilde{\pi}) \rightarrow Wh_2(A\pi)] \\ \cong \text{Ker} \left[ r_1 * \oplus r_2 *: \frac{Wh(AG)}{1 + I_1 I_2} \rightarrow Wh(A\tilde{\pi})^2 \right]; \end{aligned}$$

and in the limit

$$Wh_2^*(A\pi) \cong \varprojlim_{\alpha} \text{Ker} \left[ r_1 * \oplus r_2 *: \frac{Wh(AG)}{1 + I_1 I_2} \rightarrow Wh(A\tilde{\pi})^2 \right]. \quad (6)$$

At this point, the modified logarithm map studied in Section 2 of [10] can be applied. Let

$$I(A\tilde{\pi}) = \text{Ker}[\varepsilon: A\tilde{\pi} \rightarrow A] \quad \text{and} \quad \overline{I(A\tilde{\pi})} = H_0(\tilde{\pi}; I(A\tilde{\pi})).$$

Here,  $\varepsilon$  is the augmentation map, and  $\tilde{\pi}$  acts on  $I(A\tilde{\pi})$  by conjugation. By Theorem 2 in [10], there is a map  $\Gamma = \Gamma_{A\pi}$  and a short exact sequence

$$0 \rightarrow Wh'(A\tilde{\pi}) \xrightarrow{\Gamma_{A\pi}} \overline{I(A\tilde{\pi})} \xrightarrow{\omega} \tilde{\pi}^{ab} \rightarrow 0$$

(and similarly for  $G$ ), where  $\omega(\sum \lambda_i g_i) = \prod g_i^{\text{Tr}(\lambda_i)}$  and

$$Wh'(A\pi) = Wh(A\pi)/SK_1(A\pi).$$

The main result of Section 1 below (Theorem 1.1) says that

$$\Gamma_G(1 + I_1 I_2) = \overline{I_1 I_2},$$

where  $\overline{I_1 I_2}$  denotes the image of  $I_1 I_2$  in  $\overline{I(AG)}$ . From this we get

$$\begin{aligned} &\text{Ker} \left[ r_1 \oplus r_2: \frac{Wh'(AG)}{1 + I_1 I_2} \rightarrow Wh'(A\tilde{\pi})^2 \right] \\ &\cong \text{Ker} \left[ (\omega, r_1, r_2): H_0 \left( G; \frac{I(AG)}{I_1 I_2} \right) \rightarrow G^{ab} \oplus H_0(\tilde{\pi}; I(A\tilde{\pi}))^2 \right] \\ &\cong \text{Ker}[(\omega, i_*): H_0(G; I_{\alpha}) \rightarrow G^{ab} \oplus H_0(\tilde{\pi}; I(A\tilde{\pi}))]. \end{aligned} \quad (7)$$

Here,

$$\begin{aligned} I_\alpha &= \text{Ker}[\alpha: A\tilde{\pi} \rightarrow A\pi] \cong \text{Ker}[r_1: \mathfrak{A} \rightarrow A\tilde{\pi}] \\ &\cong \text{Ker}\left[r_1: \frac{I(AG)}{I_1 I_2} \rightarrow I(A\tilde{\pi})\right] \end{aligned}$$

(in particular, the  $G$ -action on  $I_\alpha$  factors through  $\tilde{\pi}$ ). Recall that  $r_1$  is split by the diagonal map  $\mathcal{A}: A\tilde{\pi} \rightarrow \mathfrak{A}(\subseteq (A\tilde{\pi})^2)$ .

In Proposition 3.3 and Lemma 3.4, we will see that

$$\begin{aligned} \varprojlim_{\alpha} \text{Ker}[i_*: H_0(\tilde{\pi}; I_\alpha) \rightarrow H_0(\tilde{\pi}; I(A\tilde{\pi}))] \\ \cong \varprojlim_{\alpha} \text{Coker}[\alpha_*: H_1(\tilde{\pi}; I(A\tilde{\pi})) \rightarrow H_1(\tilde{\pi}; I(A\pi))] \\ \cong H_1(\pi, A\pi) / \langle g \otimes \lambda g^n: g \in \pi, \lambda \in A, n \in \mathbb{Z} \rangle \\ = \mathcal{U}(A\pi). \end{aligned}$$

The main point here is that if  $g, h \in \pi$  and  $\langle g, h \rangle$  is not cyclic, then there is some  $\alpha: \tilde{\pi} \rightarrow \pi$  such that no liftings of  $g$  and  $h$  commute in  $\tilde{\pi}$ .

So (7) now takes the form

$$\begin{aligned} \varprojlim_{\alpha} \text{Ker}\left[r_{1*} \oplus r_{2*}: \frac{Wh'(AG)}{1 + I_1 I_2} \rightarrow Wh'(A\tilde{\pi})^2\right] \\ \cong \text{Ker}\left[\mathcal{U}(A\pi) \xrightarrow{\omega|} \varprojlim_{\alpha} \text{Ker}[G^{ab} \rightarrow \tilde{\pi}^{ab} \oplus \tilde{\pi}^{ab}] \cong H_2(\pi)\right]. \quad (8) \end{aligned}$$

That the last limit is isomorphic to  $H_2(\pi)$  follows from sequence (4). The composite of these maps is checked to be the map

$$v: \mathcal{U}(A\pi) \rightarrow H_2(\pi),$$

where  $(g \otimes \lambda h) = \text{Tr}(\lambda) \cdot (g \wedge h)$  (in the notation of Proposition 2.1 below).

It remains to see how  $SK_1(A\tilde{\pi})$  and  $SK_1(AG)$  behave in the limit. That  $\varprojlim_{\alpha} SK_1(A\tilde{\pi}) = 0$  follows easily from Lemma 22(i) in [10]. Spectral sequences show that

$$\varprojlim_{\alpha} SK_1(AG) \cong \varprojlim_{\alpha} (H_2(G)/H_2^{ab}(G)) \cong H_3(\pi)/H_3^{cv}(\pi), \quad (9)$$

where  $H_3^{cv}(\pi)$  is the subgroup of  $H_3(\pi)$  generated by induction from cyclic subgroups. So upon combining (6), (8), and (9), we get:

**THEOREM.** *For any  $p$ -group  $\pi$  and any unramified  $p$ -ring  $A$ , there is an exact sequence*

$$H_3(\pi) \xrightarrow{\eta_{A\pi}} Wh_2^*(A\pi) \rightarrow \mathcal{U}(A\pi) \xrightarrow{v} H_2(\pi).$$

Describing  $\text{Ker}(\eta_{A\pi})$  amounts to computing, for each  $\alpha: \tilde{\pi} \twoheadrightarrow \pi$ , how much of  $SK_1(AG)$  is hit by  $(1 + I_1 I_2)$ . To do this, better control is needed over  $Wh(AG)$  as an extension of  $SK_1(AG)$  by  $Wh'(AG)$ . This is obtained by regarding  $Wh(AG)$  as a quotient group of  $Wh'(A\hat{G})$  (and hence as a subquotient of  $\overline{I(AG)}$ ) for some central extension  $\hat{G} \twoheadrightarrow G$  for which  $SK_1(A\hat{G}) = 0$  (such  $G$  exists by [10, Lemma 22(i)]). The extra complications in the constructions in Sections 2 and 3 below are caused precisely by the need to work with  $\hat{G}$  instead of  $G$ .

We end this introductory section by noting that  $K_2^*(A\pi)$  can be regarded as a quotient of  $K_2^{\text{top}}(A\pi)$ , as well as of  $K_2(A\pi)$  (that  $K_2(A\pi)$  surjects onto  $K_2^*(A\pi)$  is clear once we know that  $K_2^*(A\pi)$  is finite). As usual, for any  $p$ -adic order  $\mathfrak{A}$ , we set

$$K_2^{\text{top}}(\mathfrak{A}) = \varprojlim_n K_2(\mathfrak{A}/p^n(\mathfrak{A})).$$

PROPOSITION 0.1. *For any  $p$ -group  $\pi$  and any unramified  $p$ -ring  $A$ ,*

$$K_2^*(A\pi) \cong \varprojlim \text{Coker}[K_2^{\text{top}}(A\tilde{\pi}) \rightarrow K_2^{\text{top}}(A\pi)].$$

*In particular, the natural projection  $[K_2(A\pi) \twoheadrightarrow K_2^*(A\pi)]$  factors through  $K_2^{\text{top}}(A\pi)$ .*

*Proof.* Let  $\alpha: \tilde{\pi} \twoheadrightarrow \pi$  be any surjection of a  $p$ -group  $\tilde{\pi}$  onto  $\pi$ , and define pullback rings  $\mathfrak{A}_n$  ( $n \geq 1$ ) and  $\mathfrak{A}$  from the squares

$$\begin{array}{ccc} \mathfrak{A}_n & \longrightarrow & A/p^n[\tilde{\pi}] \\ \downarrow & (1) & \downarrow \\ A/p^n[\tilde{\pi}] & \longrightarrow & A/p^n[\pi] \end{array} \quad \text{and} \quad \begin{array}{ccc} \mathfrak{A} & \longrightarrow & A\tilde{\pi} \\ \downarrow & (2) & \downarrow \\ A\tilde{\pi} & \longrightarrow & A\pi \end{array}$$

Then  $\mathfrak{A}_n \cong \mathfrak{A}/p^n\mathfrak{A}$ . So by Theorem 10 (and its proof) in [14],

$$K_1(\mathfrak{A}) \cong \varprojlim_n K_1(\mathfrak{A}_n) \quad \text{and} \quad K_1(A\tilde{\pi}) \cong \varprojlim_n K_1(A/p^n[\tilde{\pi}]).$$

The Mayer-Vietoris sequences for (1) and (2) thus induce isomorphisms

$$\begin{aligned} \text{Coker}[K_2^{\text{top}}(A\tilde{\pi}) \rightarrow K_2^{\text{top}}(A\pi)] &\cong \text{Ker}[K_1(\mathfrak{A}) \rightarrow K_1(A\tilde{\pi}) \oplus K_1(A\pi)] \\ &\cong \text{Coker}[K_2(A\tilde{\pi}) \rightarrow K_2(A\pi)]. \end{aligned}$$

Hence, in the limit,

$$K_2^*(A\pi) \cong \varprojlim_{\tilde{\pi}} \text{Coker}[K_2^{\text{top}}(A\tilde{\pi}) \rightarrow K_2^{\text{top}}(A\pi)]. \quad \blacksquare$$

## 1

We start with some general definitions. If  $G$  is any group and  $R$  any commutative ring, set

$$\overline{RG} = RG / \langle \lambda(g - hgh^{-1}) \mid \lambda \in R; g, h \in G \rangle,$$

i.e., the free  $R$ -module on the conjugacy classes of  $G$ . Let  $I(RG)$  denote the augmentation ideal, and

$$\overline{I(RG)} = I(RG) / \langle \text{conjugation} \rangle \subseteq \overline{RG}.$$

If  $\pi$  is a  $p$ -group and  $A$  an unramified  $p$ -ring, let

$$\Gamma_{A\pi}: Wh(A\pi) \rightarrow \overline{I(A\pi)}$$

be the map studied in Section 2 of [10]. The main result of this section is:

**THEOREM 1.1** *Let  $\pi$  be a  $p$ -group, and let  $\sigma, \tau \triangleleft \pi$  be normal subgroups such that  $[\sigma, \tau] = 1$ . Let  $A$  be an unramified  $p$ -ring, and set*

$$I_{A\sigma} = \text{Ker}[A\pi \rightarrow A[\pi/\sigma]] \quad \text{and} \quad I_{A\tau} = \text{Ker}[A\pi \rightarrow A[\pi/\tau]].$$

*Then*

$$\Gamma_{A\pi}(1 + I_{A\sigma}I_{A\tau}) = \overline{I_{A\sigma}I_{A\tau}},$$

where  $\overline{I_{A\sigma}I_{A\tau}}$  denotes the image of  $I_{A\sigma}I_{A\tau}$  in  $\overline{I(A\pi)}$ .

Before Theorem 1.1 can be proven, some quite technical lemmas will be needed. The first one looks rather strange, but seems to appear frequently in computations involving  $\Gamma_{A\pi}$  (see also the proof in [10] that  $\Gamma_{A\pi}$  is integer-valued).

**LEMMA 1.2.** *Let  $p$  be a prime,  $A$  a commutative ring, and  $\varphi \in \text{Aut}(A)$  a ring homomorphism such that  $\varphi(\lambda) \equiv \lambda^p \pmod{p}$  for  $\lambda \in A$ . Let  $R$  be an  $A$ -algebra,  $U \subseteq R$  a multiplicative set, and  $Y \subseteq R$  an  $A$ -submodule containing  $U$ . Set*

$$S = \left\{ \sum_i (x_i y_i - y_i x_i) \mid x_i, y_i \in R \right\}.$$

*Let  $\Phi: R \rightarrow R$  be any  $\varphi$ -linear module homomorphism such that*

$$x^{p^n} - \Phi(x^{p^{n-1}}) \in S + p^n Y \tag{1}$$



for all  $x \in U$  and  $n \geq 1$ . Then (1) holds for all  $x = \sum_i \lambda_i x_i$ , where  $\lambda_i \in A$  and  $x_i \in U$  (and all  $n \geq 1$ ).

*Proof.* Fix some  $x = \sum_i \lambda_i x_i$ , where  $\lambda_i \in A$  and  $x_i \in U$ . A typical term in  $x^{p^n}$  has the form

$$\xi = \lambda_{i_1} \cdots \lambda_{i_q} x_{i_1} \cdots x_{i_q} \quad (q = p^n).$$

Cyclic permutation of the indices  $(i_1, \dots, i_q)$  give a total of  $p^m$  terms  $\xi_1, \dots, \xi_{p^m}$  congruent to  $\xi \pmod{S}$ , where  $p^{n-m}$  is the number of cyclic permutations leaving  $(i_1, \dots, i_q)$  invariant. Then  $x_{i_1} \cdots x_{i_q}$  is a  $p^{n-m}$ th power, and the sum of these terms  $\pmod{S}$  has the form

$$\sum_{i=1}^{p^m} \xi_i \equiv \eta = p^m \bar{\lambda}^{p^{n-m}} \bar{x}^{p^{n-m}} \quad (\bar{\lambda} = \lambda_{i_1} \cdots \lambda_{i_r}, \bar{x} = x_{i_1} \cdots x_{i_r}, r = p^m).$$

If  $m = n$ , then  $\eta \in p^n Y$  by assumption. If  $m < n$ , then there are corresponding terms

$$\sum_{i=1}^{p^m} \xi'_i \equiv \eta' = p^m \varphi(\bar{\lambda})^{p^{n-m-1}} \Phi(\bar{x}^{p^{n-m-1}})$$

in the expansion of  $\Phi(x^{p^{n-1}})$ . Since  $\varphi(\bar{\lambda}) \equiv \bar{\lambda}^p \pmod{p}$  by assumption,

$$\varphi(\bar{\lambda})^{p^{n-m-1}} \equiv \bar{\lambda}^{p^{n-m}} \pmod{p^{n-m}}.$$

Also,  $\bar{x}^{p^{n-m}} \in Y$  and  $\bar{x}^{p^{n-m}} - \Phi(\bar{x}^{p^{n-m-1}}) \in S + p^{n-m} Y$  ( $\bar{x} \in U$ ), so

$$\eta - \eta' \equiv p^m [\bar{\lambda}^{p^{n-m}} \bar{x}^{p^{n-m}} - \varphi(\bar{\lambda})^{p^{n-m-1}} \Phi(\bar{x}^{p^{n-m-1}})] \in S + p^n Y.$$

The result follows by summing over all such  $(\eta, \eta')$ . ■

For Lemmas 1.3 and 1.4, we fix  $s, t \geq 1$ , and let  $F$  be the free group on generators  $h_1, \dots, h_s, k_1, \dots, k_t, g$ . Let  $\tilde{H}$  and  $\tilde{K}$  be the normal subgroups of  $F$  generated by the  $h_i$  and  $k_i$ , respectively. Set

$$G = F/[\tilde{H}, \tilde{K}], \quad H = \tilde{H}/[\tilde{H}, \tilde{K}], \quad K = \tilde{K}/[\tilde{H}, \tilde{K}],$$

and let  $H_0 = [H, G]$  and  $K_0 = [K, G]$ . Define

$$I_H = \text{Ker}[\mathbb{Z}G \rightarrow \mathbb{Z}[G/H]], \quad I_K = \text{Ker}[\mathbb{Z}G \rightarrow \mathbb{Z}[G/K]],$$

and similarly for  $I_{H_0}$  and  $I_{K_0}$ . Finally, set

$$S = \text{Ker}[\mathbb{Z}G \rightarrow \overline{\mathbb{Z}G}],$$

i.e., the subgroup of  $\mathbb{Z}G$  generated by all  $u - uvv^{-1}$  for  $u, v \in G$ .

LEMMA 1.3. For any  $u \in G$ , let  $\delta(u)$  be the unique element of the form

$$\delta(u) = h_1^{a_1} \cdots h_s^{a_s} k_1^{b_1} \cdots k_t^{b_t} g^c$$

such that  $u^{-1}\delta(u) \in [G, G]$ . Let  $\delta: \mathbb{Z}G \rightarrow \mathbb{Z}G$  be the induced ( $\hat{\mathbb{Z}}_p$ -linear) homomorphism. Then, for any

$$\begin{aligned} x &\in (S + I_H) \cap (S + I_K), \\ x &\equiv \delta(x) \pmod{I_H I_{K_0} + I_{H_0} I_K + S}. \end{aligned}$$

*Proof.* We may clearly assume  $x \in I_K$ .

Step 1. For any  $u \in G$ , let  $\delta_K(u) \in G$  be the unique element of the form

$$\delta_K(u) = h_1^{a_1} \cdots h_s^{a_s} u' \quad (u' \in \langle k_1, \dots, k_t, g \rangle)$$

such that  $u^{-1}\delta_K(u) \in H_0 = [H, G]$ . Let  $\delta_K: \mathbb{Z}G \rightarrow \mathbb{Z}G$  be the induced homomorphism. Write

$$x = \sum_i \lambda_i u_i (v_i - 1) \quad (\lambda_i \in \mathbb{Z}, u_i \in G, v_i \in K);$$

then  $\delta_K(u_i v_i) = \delta_K(u_i) \cdot v_i$  ( $v_i \in K \subseteq \langle k_1, \dots, k_t, g \rangle$ , since  $[H, K] = 1$ ). So

$$\delta_K(x) = \sum_i \lambda_i \delta_K(u_i) (v_i - 1) \equiv \sum_i \lambda_i u_i (v_i - 1) = x \pmod{I_{H_0} I_K}. \quad (1)$$

Step 2. Now write

$$\delta_K(x) = \sum_{i=1}^r \lambda_i u_i v_i, \quad (2)$$

where  $\lambda_i \in \mathbb{Z}$ ,  $u_i = h_1^{a_{i1}} \cdots h_s^{a_{is}}$ , and  $v_i \in \langle k_1, \dots, k_t, g \rangle$ . By assumption,

$$\delta_K(x) \in I_H + S = \text{Ker}[\mathbb{Z}G \rightarrow \overline{\mathbb{Z}[G/H]}].$$

In other words, since  $\langle k_1, \dots, k_t, g \rangle$  maps isomorphically onto  $G/H$ , there are elements  $\gamma_i \in \langle k_1, \dots, k_t, g \rangle$  such that

$$\sum_i \lambda_i \cdot \gamma_i v_i \gamma_i^{-1} = 0.$$

Then

$$\delta_K(x) \equiv x_1 = \sum_{i=1}^r \lambda_i (\gamma_i u_i \gamma_i^{-1}) (\gamma_i v_i \gamma_i^{-1}) \pmod{S}, \quad (3)$$

and  $x_1 \in I_H$ . Since

$$\delta(v_i) = \delta(\gamma_i v_i \gamma_i^{-1}) \equiv \gamma_i v_i \gamma_i^{-1} \pmod{K_0 = [K, G]},$$

we get in the same way as in Step 1 that

$$x_1 \equiv x_2 = \sum_{i=1}^r \lambda_i (\gamma_i u_i \gamma_i^{-1}) \delta(v_i) \pmod{I_H I_{K_0}}. \quad (4)$$

Step 3. Referring to (2), we see that

$$\delta(x) = \delta(\delta_K(x)) = \sum_i \lambda_i u_i \delta(v_i).$$

Furthermore, since  $\gamma_i \in \langle k_1, \dots, k_t, g \rangle$  and  $[H, K] = 1$ ,

$$\gamma_i u_i \gamma_i^{-1} = g^{d_i} u_i g^{-d_i}$$

for some  $d_i$ . By (1), (3), and (4),

$$x \equiv x_2 = \sum_{i=1}^r \lambda_i (g^{d_i} u_i g^{-d_i}) \delta(v_i) \pmod{I_H I_{K_0} + I_{H_0} I_K + S},$$

and it remains to show that

$$(g^d h g^{-d}) k g^c \equiv h k g^c \pmod{I_{H_0} I_K + S}$$

for any  $h \in \langle h_1, \dots, h_s \rangle$ ,  $k \in \langle k_1, \dots, k_t \rangle$ , and  $c, d \in \mathbb{Z}$ . But

$$\begin{aligned} h k g^c &= h(k g^c - g^c) + h g^c \\ &\equiv g^d h g^{-d} (k g^c - g^c) + h g^c \pmod{I_{H_0} I_K} \\ &= (g^d h g^{-d}) k g^c - g^d h g^{-d} g^c + h g^c \\ &\equiv (g^d h g^{-d}) k g^c \pmod{S}. \quad \blacksquare \end{aligned}$$

LEMMA 1.4. In the situation of Lemma 1.3, set

$$\xi = (1 - h_1) \cdots (1 - h_s) (1 - k_1) \cdots (1 - k_t) g \in \mathbb{Z}G.$$

Fix a prime  $p$ , and define  $\Phi: \mathbb{Z}G \rightarrow \mathbb{Z}G$  by

$$\Phi\left(\sum \lambda_i g_i\right) = \sum \lambda_i g_i^p \quad (\lambda_i \in \mathbb{Z}, g_i \in G).$$

Set  $H_1 = \langle [H, G], H^p \rangle \triangleleft G$ . Then for any  $n \geq 1$ ,

$$\xi^{p^n} - \Phi(\xi^{p^{n-1}}) \in S + p^n I_{H_1} I_K + p^n I_H I_{K_0} + p^n I_H^s I_K^{t+1}.$$

*Proof.* We first check that

$$\xi^{p^n} - \Phi(\xi^{p^{n-1}}) \in S + p^n \mathbb{Z}G. \quad (1)$$

Apply Lemma 1.2, with  $R = Y = \mathbb{Z}G$ ,  $U = G$ ,  $A = \mathbb{Z}$ , and  $\varphi = id$ . Note that

$$S = \left\{ \sum (x_i y_i - y_i x_i) \mid x_i, y_i \in \mathbb{Z}G \right\},$$

since the group elements generate  $\mathbb{Z}G$  additively. Since  $u^{p^n} = \Phi(u^{p^{n-1}})$  for  $u \in G$ , by definition, (1) follows by Lemma 1.2.

Now fix  $x \in \mathbb{Z}G$  such that  $\xi^{p^n} - \Phi(\xi^{p^{n-1}}) \equiv p^n x \pmod{S}$ . Then  $x$  vanishes in  $\overline{\mathbb{Z}[G/H]}$  and  $\overline{\mathbb{Z}[G/K]}$ : they are torsion free, and  $\xi \in I_H \cap I_K$ . In other words,

$$x \in (S + I_H) \cap (S + I_K),$$

and in the notation of Lemma 1.3,

$$x \equiv \delta(x) \pmod{I_H I_{K_0} + I_{H_0} I_K + S}. \quad (2)$$

Since  $\delta(S) = 0$ ,

$$\begin{aligned} \delta(x) &= \frac{1}{p^n} [\delta(\xi^{p^n}) - \delta(\Phi(\xi^{p^{n-1}}))] \\ &= \frac{1}{p^n} [(1 - h_1)^{p^n} \cdots (1 - h_s)^{p^n} \cdots (1 - k_t)^{p^n} g^{p^n} \\ &\quad - (1 - h_1^p)^{p^{n-1}} \cdots (1 - k_t^p)^{p^{n-1}} g^{p^n}]. \end{aligned} \quad (3)$$

Furthermore,

$$(1 - h_i)^{p^n} - (1 - h_i^p)^{p^{n-1}} \in p^n I_H$$

and similarly for  $k_i$  ( $p^n \mathbb{Z}G \cap I_H = p^n I_H$ , since  $\mathbb{Z}[G/H]$  is torsion free). So (3) implies that

$$\delta(x) \in I_H^s I_K^{t+1} + I_{H_1} I_K. \quad (4)$$

The lemma now follows from (2), (4), and the definition of  $x$ . ■

We now return to the situation of Theorem 1.1. For the rest of Section 1, we fix an unramified  $p$ -ring  $A$ , a  $p$ -group  $\pi$ , and subgroups  $\sigma, \tau \triangleleft \pi$  such that  $[\sigma, \tau] = 1$ . Set

$$\sigma_1 = \langle [\sigma, \pi], \text{Fr}(\sigma) \rangle \quad \text{and} \quad \tau_1 = \langle [\tau, \pi], \text{Fr}(\tau) \rangle,$$

where  $\text{Fr}(\sigma)$  and  $\text{Fr}(\tau)$  denote the Frattini subgroups. Note that  $\sigma_1, \tau_1 \triangleleft \pi$ , and  $\sigma_1 \not\subseteq \sigma$ ,  $\tau_1 \not\subseteq \tau$ . Hence, upon setting

$$X = I_{A\sigma_1} I_{A\tau} + I_{A\sigma} I_{A\tau_1} \subseteq A\pi$$

( $I_{\sigma_1} = \text{Ker}[A\pi \rightarrow A[\pi/\sigma_1]]$ , etc.), we may assume inductively that  $\Gamma_{A\pi}(1+X) = \bar{X}$  (the image of  $X$  in  $\overline{I(A\pi)}$ ).

Recall that  $\Gamma_{A\pi}$  is defined by setting

$$\Gamma_{A\pi}(x) = \log x - \frac{1}{p} \Phi(\log x) \in \overline{I(A\pi)}$$

for any  $x \in 1 + I(A\pi)$ , where  $\Phi(\sum \lambda_i g_i) = \sum \varphi(\lambda_i) g_i^p$  and  $\varphi \in \text{Gal}(A/\widehat{\mathbb{Z}}_p)$  is the Frobenius automorphism ( $\varphi(\lambda) \equiv \lambda^p \pmod{p}$ ).

LEMMA 1.5. For any  $u, v \geq 1$  and  $x \in I_{A\sigma}^u I_{A\tau}^v$ ,

$$\Gamma_{A\pi}(1+x) \equiv x \pmod{I_{A\sigma}^u I_{A\tau}^{v+1} + X \text{ in } \overline{I(A\pi)}}.$$

*Proof.* First note that

$$\begin{aligned} \Gamma(1+x) &= \left[ x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \cdots \right] - \left[ \frac{1}{p}\Phi(x) - \frac{1}{2p}\Phi(x^2) + \cdots \right] \\ &\equiv x + \sum_{k=1}^{\infty} (-1)^{k+1} \frac{1}{pk} [x^{pk} - \Phi(x^k)] \pmod{I_{A\sigma}^{2u} I_{A\tau}^{2v}}. \end{aligned}$$

(Special care is needed when  $p=2$ .) So it suffices to show that

$$x^{p^n} - \Phi(x^{p^{n-1}}) \in S' + p^n(X + I_{A\sigma}^u I_{A\tau}^{v+1}) \quad (1)$$

for all  $n \geq 1$  and  $x \in I_{A\sigma}^u I_{A\tau}^v$ , where

$$S' = \text{Ker}[I(A\pi) \rightarrow \overline{I(A\pi)}] = \left\{ \sum_i (x_i y_i - y_i x_i) \mid x_i, y_i \in A\pi \right\}.$$

We again apply Lemma 1.2, this time with  $R = A\pi$ ,

$$\begin{aligned} U &= \{ (1-h_1) \cdots (1-h_s)(1-k_1) \cdots (1-k_t) g \mid h_i \in \sigma, k_i \in \tau, \\ &\quad g \in \pi, s \geq u, t \geq v \}, \end{aligned}$$

and  $Y = X + I_{A\sigma}^u I_{A\tau}^{v+1}$ . Since  $[\sigma, \tau] = 1$ ,  $U$  is multiplicative. Relation (1) holds for all  $x \in U$  by Lemma 1.4. So by Lemma 1.2, (1) holds for all

$$x \in I_{A\sigma}^u I_{A\tau}^v = \left\{ \sum_i \lambda_i x_i \mid \lambda_i \in A, x_i \in U \right\}. \quad \blacksquare$$

*Poof of Theorem 1.1.* First note that  $\Gamma(1 + I_{A\sigma} I_{A\tau}) \subseteq \overline{I_{A\sigma} I_{A\tau}}$  by Lemma 1.5. As remarked,  $\Gamma(1+X) = \bar{X}$  by induction, and hence

$$\bar{X} \subseteq \Gamma(1 + I_{A\sigma} A_{A\tau}).$$

We next check that  $\Phi(x) \in X$  for all  $x \in I_{A\sigma}I_{A\tau}$ . It suffices to show this when  $x = (1-h)(1-k)g$  for any  $h \in \sigma$ ,  $k \in \tau$ , and  $g \in \pi$ . For such  $x$ ,

$$\Phi(x) = g^p - (hg)^p - (kg)^p + (hkg)^p = (1-h')(1-k')g,$$

where

$$h' = h(ghg^{-1}) \cdots (g^{p-1}hg^{-p+1}) \quad \text{and} \quad k' = k(gkg^{-1}) \cdots (g^{p-1}kg^{-p+1})$$

(recall that  $[\sigma, \tau] = 1$ ). Furthermore,

$$h' \in \sigma_1 = \langle [\sigma, \pi], \text{Fr}(\sigma) \rangle \quad \text{and} \quad k' \in \tau_1 = \langle [\tau, \pi], \text{Fr}(\tau) \rangle,$$

and so  $\Phi(x) \in I_{A\sigma_1}I_{A\tau_1} \subseteq X$ .

Thus, for any  $n \geq 2$  and any  $x \in I_{A\sigma}I_{A\tau}$ ,

$$\begin{aligned} \Gamma(1 + p^n x) &\equiv \log(1 + p^n x) \pmod{\bar{X}} \\ &\equiv p^n x \pmod{\bar{X} + p^{n+1}\overline{I_{A\sigma}I_{A\tau}}}. \end{aligned}$$

By taking successive approximations, it follows that

$$\Gamma(1 + I_{A\sigma}I_{A\tau}) \supseteq p^2 I_{A\sigma}I_{A\tau}.$$

Finally, note that for  $t$  large enough,

$$I'_{A\tau} \subseteq p^2 I_{A\tau} \quad \text{and} \quad I_{A\sigma} I'_{A\tau} \subseteq p^2 I_{A\sigma} I_{A\tau}.$$

So by Lemma 1.5 and downward induction,

$$\Gamma(1 + I_{A\sigma}I_{A\tau}) \supseteq I_{A\sigma} I'_{A\tau}$$

for all  $t \geq 1$ . In particular,  $\Gamma(1 + I_{A\sigma}I_{A\tau}) = \overline{I_{A\sigma}I_{A\tau}}$ . ■

## 2

Throughout this section,  $A$  will be a fixed unramified  $p$ -ring (for some fixed prime  $p$ ), and  $\varphi \in \text{Gal}(A/\mathbb{Z}_p)$  the Frobenius automorphism. We let  $\text{Tr} = \text{Tr}_{A/\mathbb{Z}_p}$  denote the trace map; it is surjective by Proposition VIII.1.3 in [15]. The goal is to describe  $K_1(A\tilde{\pi}, I)$  when  $\tilde{\pi}$  is a  $p$ -group and  $I = \text{Ker}[A\tilde{\pi} \rightarrow A\pi]$  for some surjection  $\tilde{\pi} \rightarrow \pi$ .

We first list some basic facts involving  $H_2(\pi)$  which will be needed frequently.

PROPOSITION 2.1. (i) For any extension  $1 \rightarrow \rho \rightarrow \pi \xrightarrow{\alpha} \pi \rightarrow 1$  of groups, there is a 5-term homology exact sequence natural in  $\alpha$ :

$$H_2(\tilde{\pi}) \xrightarrow{\alpha_*} H_2(\pi) \xrightarrow{\delta^\alpha} \rho/[\tilde{\pi}, \rho] \rightarrow \tilde{\pi}^{ab} \rightarrow \pi^{ab} \rightarrow 0.$$

(ii) If  $1 \rightarrow R \rightarrow F \xrightarrow{\alpha} \pi \rightarrow 1$  is an extension, where  $F$  is free and  $\exp H_2(\pi) \mid q < \infty$ , then the composite

$$H_2(\pi) \xrightarrow{\delta^\alpha} R/[F, R] \rightarrow (R/[F, R]) \otimes \mathbb{Z}/q$$

is injective.

(iii) If  $\pi = F/R$  where  $F$  is free, then  $H_2(\pi) \cong (R \cap [F, F])/[R, F]$ . In particular, if  $g, h \in \pi$  and  $gh = hg$ , then they determine an element  $g \wedge h \in H_2(\pi)$ :  $g \wedge h = [\tilde{g}, \tilde{h}] \in R \cap [F, F]$  for any liftings  $\tilde{g}, \tilde{h} \in F$  of  $g$  and  $h$ . We define

$$H_2^{ab} = \langle g \wedge h \in H_2(\pi) : g, h \in \pi, gh = hg \rangle \subseteq H_2(\pi).$$

(iv) For any  $p$ -group  $\pi$ ,

$$SK_1(A\pi) \cong H_2(\pi)/H_2^{ab}(\pi).$$

For any central extension  $1 \rightarrow \rho \rightarrow \tilde{\pi} \xrightarrow{\alpha} \pi \rightarrow 1$  of  $p$ -groups,

$$SK_1(A\tilde{\pi}) \cong \text{Ker}(\delta^\alpha)/\langle g \wedge h : g, h \in \pi, gh = hg, g \wedge h \in \text{Ker}(\delta^\alpha) \rangle.$$

In particular,  $SK_1(A\tilde{\pi}) \cong SK_1(A\pi)$  if  $\delta^\alpha = 0$ , and  $SK_1(A\tilde{\pi}) = 0$  if  $\delta^\alpha$  is injective.

*Proof.* The homology sequence is constructed, e.g., in [6, Corollary VI.8.2]. To see (ii), note that  $\text{Coker}(\delta^\alpha) \subseteq F^{ab}$  is free abelian and  $H_2(F) = 0$ , so that  $\delta^\alpha$  is split injective. For (iv), see Theorem 3 and Lemma 22(i) in [10]. ■

From now on, for any extension  $1 \rightarrow \rho \rightarrow \tilde{\pi} \xrightarrow{\alpha} \pi \rightarrow 1$  of groups, we set

$$I_{A\alpha} = \text{Ker}[A\tilde{\pi} \xrightarrow{\alpha} A\pi].$$

Maps

$$\omega^* = \omega_{A\alpha}^* : I_{A\alpha} \rightarrow A \otimes \rho^{ab} \quad \text{and} \quad \omega = \omega_{A\alpha} : I_{A\alpha} \rightarrow \hat{\mathbb{Z}}_p \otimes \rho^{ab}$$

are defined by setting

$$\omega^* \left( \sum_i \lambda_i (h_i - 1) g_i \right) = \sum_i \lambda_i \otimes h_i \quad (\lambda_i \in A, h_i \in \rho, g_i \in \tilde{\pi}),$$

and letting  $\omega = (\text{Tr} \otimes 1) \omega^*$ . Note that  $\hat{\mathbb{Z}}_p \otimes \rho^{ab} \cong \rho^{ab}$  if  $\rho$  is a  $p$ -group.

PROPOSITION 2.2. Let  $1 \rightarrow \rho \rightarrow \tilde{\pi} \xrightarrow{\alpha} \pi \rightarrow 1$  be an extension of groups.

(i)  $\text{Ker}(\omega_{A\alpha}^*) = I_{A\alpha}I$ , where  $I = I(A\tilde{\pi})$ . In other words,  $\omega_{A\alpha}^*$  induces an isomorphism.

$$I_{A\alpha}/I_{A\alpha}I \cong A \otimes \rho^{ab}.$$

(ii) If  $\tilde{\pi}$  is a  $p$ -group and  $\rho \subseteq Z(\tilde{\pi})$  is central, then

$$\Gamma_{A\tilde{\pi}}(1 + I_{A\alpha}) = \text{Ker}[\omega_{A\alpha}: \bar{I}_{A\alpha} \rightarrow \rho] \subseteq \overline{I(A\tilde{\pi})}.$$

*Proof.* (i) Define

$$s: A \otimes \rho^{ab} \rightarrow I_{A\alpha}/I_{A\alpha}I$$

by setting  $s(\lambda \otimes h) = \lambda(h - 1)$ . This is clearly a homomorphism:

$$s(\lambda \otimes hk) - s(\lambda \otimes h) - s(\lambda \otimes k) = \lambda(1 - h)(1 - k) \in (I_{A\alpha})^2$$

for any  $\lambda \in A$  and  $h, k \in p$ . That  $\omega_{A\alpha}^*$  and  $s$  are inverses is easy.

(ii) By (i),  $I_{A\alpha}$  is generated by  $I_{A\alpha}I$  and  $I(A\rho)$  (the augmentation ideal in  $A\rho$ ). Hence, by Theorem 1.1 (applied to the pair  $(\sigma, \tau) = (\rho, \tilde{\pi})$ ) and Theorem 2 in [10],

$$\begin{aligned} \Gamma(1 + I_{A\alpha}) &= \Gamma(1 + I_{A\alpha}I) + \Gamma(1 + I(A\rho)) = \overline{I_{A\alpha}I} + \text{Ker}[\omega_{A\alpha}: I(A\rho) \rightarrow \rho] \\ &= \text{Ker}[\omega: \bar{I}_{A\alpha} \rightarrow \rho]. \quad \blacksquare \end{aligned}$$

This can be applied to give the following description of  $K_1(A\pi)$ :

PROPOSITION 2.3. Let  $\pi$  be a  $p$ -group, and set  $I = I(A\pi)$ . Then

$$K_1(A\pi) \cong A^* \oplus K_1(A\pi, I)$$

(canonically), and there is a natural exact sequence

$$0 \rightarrow K_1(A\pi, I) \xrightarrow{\mu} (A \otimes \pi^{ab}) \oplus \text{Wh}(A\pi) \xrightarrow{\nu} A \otimes \pi^{ab}.$$

Here,  $\mu(1 + x) = (\omega_{A\pi}^*(x), [1 + x])$  for  $x \in I$ , and

$$\begin{aligned} \nu(\lambda \otimes g, [1 + x]) &= ((1 - \varphi) \otimes 1)(\lambda \otimes g - \omega_{A\pi}^*(x)) \\ &= (\lambda - \varphi(\lambda)) \otimes g - \omega_{A\pi}^* \Gamma_{A\pi}(1 + x). \end{aligned}$$

By  $\omega_{A\pi}^*: I(A\pi) \rightarrow A \otimes \pi^{ab}$  is meant  $\omega_{A\alpha}^*$  when  $\alpha: \pi \rightarrow 1$ .

*Proof.* We first check that the two formulas for  $\nu$  agree, i.e., that

$$\omega_{A\pi}^* \Gamma_{A\pi}(1 + x) = ((1 - \varphi) \otimes 1)(\omega_{A\pi}^*(x)) \in A \otimes \pi^{ab}$$



for any  $x \in I$ . It clearly suffices to do this when  $\pi$  is abelian, and by Proposition 2.2 it suffices to consider the cases (a)  $x \in I^2$  and (b)  $x = \lambda(g-1)$  ( $\lambda \in A, g \in \pi$ ). In case (a),

$$\omega_{A\pi}^*(x) = 0 \quad \text{and} \quad \Gamma_{A\pi}(1+x) \in I^2 = \text{Ker}(\omega_{A\pi}^*)$$

by Proposition 2.2 and Theorem 1.1. In case (b),

$$\begin{aligned} & \Gamma(1 + \lambda(g-1)) \\ &= \frac{1}{p} \log \left( \frac{(1 + \lambda(g-1))^p}{1 + \varphi(\lambda)(g^p - 1)} \right) \\ &\equiv \frac{1}{p} \log \left( \frac{1 + \lambda^p(g^p - 1) + p(\lambda - \lambda^p)(g-1)}{1 + \varphi(\lambda)(g^p - 1)} \right) \pmod{I_2} \\ &\equiv \frac{1}{p} \log(1 + (\lambda^p - \varphi(\lambda))(g^p - 1) + p(\lambda - \lambda^p)(g-1)) \\ &\equiv \frac{1}{p} (\lambda^p - \varphi(\lambda))(g^p - 1) + (\lambda - \lambda^p)(g-1) \pmod{I^2}, \end{aligned}$$

and so

$$\begin{aligned} \omega_{A\pi}^* \Gamma_{A\pi}(1 + \lambda(g-1)) &= \frac{1}{p} (\lambda^p - \varphi(\lambda)) \otimes g^p + (\lambda - \lambda^p) \otimes g \\ &= (\lambda - \varphi(\lambda)) \otimes g. \end{aligned}$$

Since  $v(Wh(A\pi)) \subseteq v(A \otimes \pi^{ab})$ ,

$$\text{Ker}(v) = \langle (1 \otimes g, 1), (\omega_{A\pi}^*(x), [1+x]): g \in \pi, x \in I(A\pi) \rangle = \text{Im}(\mu).$$

Furthermore,

$$\text{Ker}(\mu) \subseteq \text{Ker}[K_1(A\pi, I) \rightarrow Wh(A\pi)] = \pi^{ab},$$

but  $\mu(g) = (1 \otimes g, 1)$  for  $g \in \pi$ , and so  $\mu$  is injective.

The isomorphism  $K_1(A\pi) \cong K_1(A\pi, I) \oplus A^*$  is induced by the inclusion  $A \subseteq A\pi$  and the augmentation map  $A\pi \rightarrow A$ . ■

One more technical lemma will be needed:

LEMMA 2.4. *Let  $G$  be any group,  $H, K \triangleleft G$  normal subgroups such that  $H \cap K = 1$ , and set*

$$I_H = \text{Ker}[AG \rightarrow A[G/H]] \quad \text{and} \quad I_K = \text{Ker}[AG \rightarrow A[G/K]].$$

Then  $I_H \cap I_K = I_H I_K$ . Furthermore, if  $S \subseteq G$  is a set of coset representatives for  $H \times K \subseteq G$ , then  $I_H \cap I_K$  is a free  $A$ -module with basis

$$X = \{(h-1)(k-1)g : h \in H-1, k \in K-1, g \in S\}.$$

*Proof.* Clearly,  $AX \subseteq I_H I_K \subseteq I_H \cap I_K$ . Each element of  $G - (HS \cup KS)$  (i.e., each element of the form  $hkg$  for  $h \in H-1, k \in K-1, g \in S$ ) occurs as a term in exactly one element of  $X$ , and each element of  $X$  has exactly one such term. This implies first that the elements of  $X$  are  $A$ -linearly independent, and second that any  $x \in AG$  is congruent (mod  $AX$ ) to some  $y = \sum \lambda_i g_i$  for  $\lambda_i \in A$  and  $g_i \in HS \cup KS$ . In particular, if  $x \in I_H \cap I_K$ , then  $y = 0$ . In other words,  $I_H \cap I_K \subseteq AX$ , and we are done. ■

**THEOREM 2.5.** Let  $1 \rightarrow \rho \rightarrow \tilde{\pi} \xrightarrow{\alpha} \pi \rightarrow 1$  be an extension of  $p$ -groups, set

$$I_{A\alpha} = \text{Ker}[A\tilde{\pi} \rightarrow A\pi] \quad \text{and} \quad M_{A\alpha} = \text{Ker}[\omega_{A\alpha}: H_0(\rho; I_{A\alpha}) \rightarrow \rho^{ab}],$$

and let  $\iota: M_{A\alpha} \rightarrow H_0(\rho; I_{A\alpha})$  be the inclusion. Then there is an exact sequence

$$\begin{aligned} SK_1(A\rho) &\xrightarrow{s} K_1(A\tilde{\pi}, I_{A\alpha}) \xrightarrow{\Psi} A \otimes (\rho/[\tilde{\pi}, \rho]) \oplus H_0(\pi; M_{A\alpha}) \\ &\xrightarrow{(1-\varphi) \otimes 1 - (\omega_{A\alpha}^*)/\pi} A \otimes (\rho/[\tilde{\pi}, \rho]) \end{aligned}$$

which is natural with respect to maps between extensions. Here,

- (i)  $s$  is the composite  $SK_1(A\rho) \subseteq K_1(A\rho, I(A\rho)) \rightarrow K_1(A\tilde{\pi}, I_{A\alpha})$ , and
- (ii)  $\Psi = (\Psi_1, \Psi_2)$ , where  $\Psi_1(1+x) = \omega_{A\alpha}^*(x)$ .

Furthermore, the composite

$$\begin{aligned} H_2(\pi) &\xrightarrow{\lambda_\pi} K_2(A\pi)/\{-1, \pi\} \xrightarrow{\partial} K_1(A\tilde{\pi}, I_{A\alpha}) \\ &\xrightarrow{\Psi} A \otimes (\rho/[\tilde{\pi}, \rho]) \oplus H_0(\pi; M_{A\alpha}) \end{aligned}$$

(where  $\lambda_\pi$  denotes the map  $\lambda''(\pi)$  defined in Section 4.3 of [8]) is equal to the pair  $(\delta^\alpha, 0)$ .

*Proof.* Define a group  $G$  and order  $\mathfrak{A}$  as pullback:

$$\begin{array}{ccc} G & \xrightarrow{r_2} & \tilde{\pi} \\ \downarrow r_1 & & \downarrow \alpha \\ \tilde{\pi} & \xrightarrow{\alpha} & \pi \end{array} \quad \text{and} \quad \begin{array}{ccc} \mathfrak{A} & \xrightarrow{r_2} & A\tilde{\pi} \\ \downarrow r_1 & & \downarrow \\ A\tilde{\pi} & \longrightarrow & A\pi \end{array}$$

Let  $\Delta: \tilde{\pi} \rightarrow G$  be the diagonal map (regarding  $G$  as a subgroup of  $\tilde{\pi} \times \tilde{\pi}$ ). Set  $\rho_i = \text{Ker}(r_i) \subseteq G$ , and note that  $\rho_1 \cong \rho_2 \cong \rho$  and  $G/(\rho_1 \times \rho_2) \cong \pi$ . Set

$$I_1 = \text{Ker}[r_1: AG \rightarrow A\tilde{\pi}] \quad \text{and} \quad I_2 = \text{Ker}[r_2: AG \rightarrow A\tilde{\pi}].$$

Then  $\mathfrak{U} \cong AG/(I_1 \cap I_2)$  (since  $\mathfrak{U} \subseteq A\tilde{\pi} \times A\tilde{\pi}$ ).

*Step 1.* By Proposition 2.3, there is an exact sequence

$$0 \rightarrow K_1(AG) \rightarrow A^* \oplus (A \otimes G^{ab}) \oplus Wh(AG) \xrightarrow[-\omega_{AG}^* \Gamma_{AG}]{(1-\varphi) \otimes 1} A \otimes G^{ab}. \quad (1)$$

Furthermore,  $K_1(\mathfrak{U}) \cong K_1(AG)/(1 + I_1 I_2)$  ( $I_1 \cap I_2 = I_1 I_2$  by Lemma 2.4), and

$$I_1 I_2 \subseteq I(AG)^2 = \text{Ker}(\omega_{AG}^*)$$

by Lemma 2.2. So upon dividing (1) out by  $1 + I_1 I_2$ , we get the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \rightarrow & K_1(\mathfrak{U}) & \rightarrow & A^* \oplus (A \otimes G^{ab}) \oplus \frac{Wh(AG)}{1 + I_1 I_2} & \rightarrow & A \otimes G^{ab} \\ & & \downarrow K_1(r_1) & & \downarrow (1, 1 \otimes r_1^{ab}, Wh(r_1)) & & \downarrow 1 \otimes r_1^{ab} \\ 0 & \rightarrow & K_1(A\tilde{\pi}) & \rightarrow & A^* \oplus (A \otimes \tilde{\pi}^{ab}) \oplus Wh(A\tilde{\pi}) & \rightarrow & A \otimes \tilde{\pi}^{ab} \end{array} \quad (2)$$

For convenience, we set  $Wh(\mathfrak{U}) = Wh(AG)/(1 + I_1 I_2)$ .

Now,  $K_1(A\tilde{\pi}, I_{A\alpha}) = \text{Ker}(K_1(r_1))$  by definition [9]. Furthermore, since  $r_1$  is split by  $\Delta$ ,

$$\text{Ker}(r_1^{ab}) = \rho_1/[G, \rho_1] = H_0(G; \rho_1^{ab}) \cong H_0(\pi; \rho^{ab}) = \rho/[\tilde{\pi}, \rho].$$

So the kernels in (2) form an exact sequence

$$\begin{aligned} 0 \rightarrow K_1(A\tilde{\pi}, I_{A\alpha}) &\xrightarrow{\psi'} A \otimes (\rho/[\tilde{\pi}, \rho]) \oplus \text{Ker}(Wh_1(r_1)) \\ &\xrightarrow[-\omega_{AG}^* \Gamma_{AG}]{(1-\varphi) \otimes 1} A \otimes \rho/[\tilde{\pi}, \rho]. \end{aligned} \quad (3)$$

*Step 2.* Now let  $1 \rightarrow \sigma \rightarrow \hat{G} \xrightarrow{\beta} G \rightarrow 1$  be any central extension such that

- (a)  $\text{Ker}(\delta^\beta) = \text{Im}[H_2(\rho_1 \times \rho_2) \oplus H_2(\tilde{\pi}) \xrightarrow{\text{incl} + \Delta^*} H_2(G)]$ , and
- (b)  $\Delta$  lifts to a map  $\hat{\Delta}: \tilde{\pi} \rightarrow \hat{G}$ .

For example,  $(\hat{G}, \beta)$  can be constructed by first choosing any central extension

$$1 \rightarrow \sigma \rightarrow G' \xrightarrow{\beta'} G \rightarrow 1$$

satisfying (a) (see Proposition V.5.1 in [12]), and then setting

$$[\hat{G}, \beta] = [G', \beta'] - r_2^* \mathcal{A}^* [G', \beta'] \in H^2(G, \sigma).$$

Then  $\delta^\beta = \delta^{\beta'}$  ( $\delta^{\beta'} \circ \mathcal{A}_* = 0$  by (a)), and  $\mathcal{A}^* [\hat{G}, \beta] = 0$ .

Set  $\hat{\rho}_i = \beta^{-1}(\rho_i)$  ( $i = 1, 2$ ), and note that  $[\hat{\rho}_1, \hat{\rho}_2] = 1$  by construction. Furthermore, by (a) and Proposition 2.1(iv),

$$\text{incl} + \mathcal{A}_*: SK_1(A\hat{\rho}_1) \oplus SK_1(A\hat{\pi}) \rightarrow SK_1(A\hat{G}) \quad (4)$$

is onto, and  $SK_1(A\hat{\rho}_1) \cong SK_1(A\rho_1)$ . Define ideals

$$J = I(A\hat{G}), \quad J_0 = \text{Ker}[\beta: A\hat{G} \rightarrow AG],$$

$$J_i = \text{Ker}[r_i\beta: A\hat{G} \rightarrow A\hat{\pi}] \quad (i = 1, 2).$$

Then  $AG = A\hat{G}/J_0$  and  $I_i = J_i/J_0$  ( $i = 1, 2$ ), and so

$$Wh(\mathfrak{A}) = Wh(AG)/(1 + I_1 I_2) \cong Wh(A\hat{G})/(1 + J_0 + J_1 J_2).$$

By Theorem 1.1 and Proposition 2.2(ii),

$$\Gamma(1 + J_0 + J_1 J_2) = \text{Ker}[\omega_{A\beta}: J_0 \rightarrow \sigma] + J_1 J_2$$

(or rather, its image in  $\overline{I(A\hat{G})}$ ).

Now, identify  $I(A\hat{G})$  with  $H_0(G; I(A\hat{G}))$  ( $G$  acting by conjugation) and write  $K = \text{Ker}(\omega_{A\beta})$  for short. Then Theorem 2 in [10] applies to give the following diagram with exact rows:

$$\begin{array}{ccccccc} SK_1(A\hat{G}) & \longrightarrow & Wh(\mathfrak{A}) & \xrightarrow{\Gamma_{A\hat{G}}} & H_0(G; J/(K + J_1 J_2)) & \xrightarrow{\omega_{A\hat{G}}} & \hat{G}^{ab} \longrightarrow \\ \downarrow SK_1(r_1\beta) & & \downarrow Wh(r_1) & & \downarrow \overline{I(r_1)} & & \downarrow r_1^{ab} \\ 0 \longrightarrow & SK_1(A\hat{\pi}) & \longrightarrow & Wh(A\hat{\pi}) & \xrightarrow{\Gamma_{A\hat{\pi}}} & H_0(G; I(A\hat{\pi})) & \xrightarrow{\omega_{A\hat{\pi}}} \hat{\pi}^{ab} \longrightarrow \end{array}$$

As in (2), the vertical maps are all split by maps induced by  $\mathcal{A}$  or  $\hat{\mathcal{A}}$ , and so the kernels form an exact sequence. By (4),  $\text{Ker}(SK_1(r_1\beta))$  is generated by  $SK_1(A\hat{\rho}_1) \cong SK_1(A\rho_1)$ . It follows that  $\Gamma_{A\hat{G}}$  induces an isomorphism (where  $\hat{r}_1 = r_1\beta: \hat{G} \rightarrow \hat{\pi}$ )

$$\begin{aligned} & \text{Ker} \left[ \frac{Wh(\mathfrak{A})}{SK_1(A\rho_1)} \xrightarrow{r_{1*}} Wh(A\hat{\pi}) \right] \\ & \cong \text{Ker} \left[ \omega_{A\hat{r}_1}: H_0 \left( G; \frac{J_1}{K + J_1 J_2} \right) \rightarrow \hat{\rho}_1 / [\hat{G}, \hat{\rho}_1] \right]. \end{aligned} \quad (5)$$

Step 3. Now consider the group

$$\begin{aligned} \text{Ker} \left[ r_2: \frac{J_1}{K + J_1 J_2} \rightarrow I_{A\alpha} \right] &= \frac{J_1 \cap J_2}{K + J_1 J_2} = \frac{J_0 + J_1 J_2}{K + J_1 J_2} \\ &\cong \frac{J_0}{K + (J_0 \cap J_1 J_2)}. \end{aligned}$$

Using Lemma 2.4 (the fact that  $X$  is a basis for  $I_1 I_2 = J_1 J_2 / J_0$ ), we see that  $J_0 \cap J_1 J_2$  is generated by elements

$$\begin{aligned} (1-h)(1-k)g - (1-h')(1-k')g \\ = (h' - h)(1-k)g + (1-h')(k' - k)g \in JJ_0 \end{aligned}$$

(where  $h \in \hat{\rho}_1$ ,  $k \in \hat{\rho}_2$ ,  $g \in \hat{G}$ , and  $h^{-1}h'$ ,  $k^{-1}k' \in \sigma$ ). Since  $JJ_0 \subseteq K$ ,

$$\text{Ker}[r_2: J_1/(K + J_1 J_2) \rightarrow I_{A\alpha}] \cong J_0/K = J_0/\text{Ker}(\omega_{A\beta}) \cong \sigma.$$

In other words, there is a short exact sequence

$$0 \rightarrow \sigma \xrightarrow{f} J_1/(K + J_1 J_2) \xrightarrow{r_2} I_{A\alpha} \rightarrow 0, \quad (6)$$

where  $f(g) = \lambda(g - 1)$  for any  $g \in \sigma$ , and any  $\lambda \in A$  such that  $\text{Tr}(\lambda) = 1$ .

Step 4. Write  $\hat{\rho} = \hat{\rho}_1 = \text{Ker}[\hat{\rho}_1: G \rightarrow \tilde{\pi}]$ , and consider the maps

$$\omega_{A\alpha}: I_{A\alpha} \rightarrow \rho^{ab} \quad \text{and} \quad \omega_{A\hat{\rho}_1}: J_1 \rightarrow \hat{\rho}^{ab}$$

( $J_1 = \text{Ker}[\hat{\rho}_1: A\hat{G} \rightarrow A\tilde{\pi}]$  defined as before). Since

$$K + J_1 J_2 = \text{Ker}(\omega_{A\beta}) + J_1 J_2 \subseteq \text{Ker}(\hat{\omega}),$$

there is a commutative diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & \sigma & \xrightarrow{f} & H_0(\rho_1 \times \rho_2; J_1/(K + J_1 J_2)) & \xrightarrow{r_2} & H_0(\rho; I_{A\alpha}) \rightarrow 0 \\ & & \downarrow \text{id} & & \downarrow \hat{\omega} & & \downarrow \omega \\ 0 & \rightarrow & \sigma & \longrightarrow & \hat{\rho}^{ab} & \xrightarrow{r_2} & \rho^{ab} \rightarrow 0 \end{array} \quad (7)$$

Both rows are right exact: the top row by (6) and the bottom row by the homology sequence for the extension

$$1 \rightarrow \sigma \rightarrow \hat{\rho} \xrightarrow{\beta_2} \rho \rightarrow 1$$

where  $\beta_2 = (r_2\beta) \mid \hat{\rho}$ . But  $\delta^{\beta_2} = 0$  by construction (see (a)), and so the rows in (7) are also exact at  $\sigma$ .

It follows that

$$\text{Ker}(\hat{\omega}) \cong \text{Ker}(\omega) = M_{A\alpha}.$$

Hence, there is an exact sequence

$$\text{Coker}(H_1(\pi; \hat{\omega})) \rightarrow H_0(\pi; M_{A\alpha}) \rightarrow H_0(G; J_1/(K + J_1J_2)) \rightarrow H_0(\pi; \hat{\rho}^{ab}). \quad (8)$$

The homology spectral sequences for  $\tilde{\pi} \rightarrow^\alpha \pi$  and  $G \rightarrow^{\bar{\alpha}} \pi$  induce a commutative diagram

$$\begin{array}{ccccccc} H_3(\pi) & \xrightarrow{\partial} & H_1(\pi; \rho^{ab}) & \rightarrow & H_2(\tilde{\pi})/H_2(\rho) & \rightarrow & H_2(\pi) \\ \downarrow \text{id} & & \downarrow \text{diag} & & \downarrow \Delta_* & & \downarrow \text{id} \\ H_3(\pi) & \xrightarrow{\partial} & H_1(\pi; \rho_1^{ab} \times \rho_2^{ab}) & \rightarrow & H_2(G)/H_2(\rho_1 \times \rho_2) & \rightarrow & H_2(\pi) \end{array}$$

where the vertical maps are split injective (split by  $r_2$ ). So

$$H_1(\pi; \rho^{ab}) \cong \text{Coker}[\text{incl} + \Delta_*: H_2(\rho_1 \times \rho_2) \oplus H_2(\tilde{\pi}) \rightarrow H_2(G)],$$

and by (a) in Step 2,

$$H_1(\pi; \rho^{ab}) \cong \text{Im}[\delta^\beta: H_2(G) \rightarrow \sigma]. \quad (9)$$

On the other hand, if  $\partial: H_1(\pi; \rho^{ab}) \rightarrow \sigma$  is the boundary map induced by the  $\pi$ -action on the bottom row of (7), then

$$\text{Im}(\partial) = \text{Ker}[\sigma \rightarrow H_0(\pi; \hat{\rho}^{ab})] = \text{Ker}[\sigma \rightarrow \hat{G}^{ab}] = \text{Im}(\delta^\beta); \quad (10)$$

$H_0(\pi; \hat{\rho}^{ab})$  injects into  $\hat{G}^{ab}$  since  $\hat{G} \rightarrow \tilde{\pi}$  is split by  $\hat{A}$ . So by (9) and (10),  $\partial$  is injective.

Referring again to diagram (7), this says that

$$\text{Coker}[H_1(\pi; \sigma) \rightarrow H_1(\pi; \hat{\rho}^{ab})] \cong \text{Ker}[\partial: H_1(\pi; \rho^{ab}) \rightarrow \sigma] = 0.$$

So  $H_1(\pi; \hat{\omega})$  is onto, and (8) takes the form

$$H_0(\pi; M_{A\alpha}) \cong \text{Ker}[\hat{\omega}: H_0(G; J_1/(K + J_1J_2)) \rightarrow H_0(\pi; \hat{\rho}_1^{ab})]. \quad (11)$$

Combining this with (5) and (3) gives, finally, the exact sequence

$$\begin{aligned} 0 \rightarrow K_1(A\tilde{\pi}, I_{A\alpha})/SK_1(A\rho) &\xrightarrow{\Psi} A \otimes (\rho/[\tilde{\pi}, \rho]) \oplus H_0(\pi; M_{A\alpha}) \\ &\rightarrow A \otimes (\rho/[\tilde{\pi}, \rho]). \end{aligned} \quad (12)$$

The descriptions of the maps in (12) follow directly from the construc-

tions. That  $\Psi$  is uniquely defined (independently of the choice of  $\widehat{G}$ ) and natural follows by taking pullbacks: for example, if  $\widehat{G}_1$  and  $\widehat{G}_2$  are two central extensions of  $G$  satisfying conditions (a) and (b) in Step 2, then their pullback over  $G$  also satisfies (a) and (b).

Finally, the description of  $\Psi \partial\lambda_\pi$  follows from the construction of (3) and the commutativity of the square:

$$\begin{array}{ccc} H_2(\pi) & \xrightarrow{\delta^\pi} & \rho/[\tilde{\pi}, \rho] \\ \downarrow \lambda_\pi & & \downarrow \\ K_2(A\pi)/\{-1, \pi\} & \xrightarrow{\partial} & K_1(A\tilde{\pi}, I_{A\alpha}) \quad \blacksquare \end{array}$$

A description of the map  $\Psi_2: K_1(A\tilde{\pi}, I_{A\alpha}) \rightarrow H_0(\pi; M_{A\alpha})$  in Theorem 2.5 can be extracted from the proof of the theorem, but is technically quite difficult to write down. The composite

$$(\iota/\pi) \circ \Psi_2: K_1(A\tilde{\pi}, I_{A\alpha}) \rightarrow H_0(\tilde{\pi}; I_{A\alpha})$$

is, on the other hand, much more natural, and is basically a relative version of the modified logarithm map  $\Gamma_{A\tilde{\pi}}$ . More precisely:

**PROPOSITION 2.6.** *Let  $1 \rightarrow \rho \rightarrow \tilde{\pi} \xrightarrow{\alpha} \pi \rightarrow 1$  be any extension of  $p$ -groups, let  $G \subseteq \tilde{\pi} \times \tilde{\pi}$  and  $\mathfrak{A} \subseteq A\tilde{\pi} \times A\tilde{\pi}$  be the pullbacks over  $\pi$  and  $A\pi$ , and let  $r_1$  and  $r_2$  be the projections of  $\mathfrak{A}$  onto  $A\tilde{\pi}$ . Let  $\gamma: AG \rightarrow \mathfrak{A}$  be the projection, and define*

$$\Gamma_{A\alpha}: K_1(A\tilde{\pi}, I_{A\alpha}) \rightarrow H_0(\tilde{\pi}; I_{A\alpha}) \quad (I_{A\alpha} = \text{Ker}[A\tilde{\pi} \rightarrow A\pi])$$

to be the composite

$$\begin{aligned} K_1(A\tilde{\pi}, I_{A\alpha}) &\xrightarrow{r_{2*}^{-1}} \text{Ker}[r_{1*}: K_1(\mathfrak{A}) \rightarrow K_1(A\tilde{\pi})] \\ &\leftarrow \text{Ker}[r_1\gamma: K_1(AG) \rightarrow K_1(A\tilde{\pi})] \\ &\xrightarrow{\Gamma_{AG}} \text{Ker}[r_1\gamma: \overline{I(AG)} \rightarrow \overline{I(A\tilde{\pi})}] \cong H_0(G; I_{A\alpha}) \\ &\xrightarrow{r_2} H_0(\tilde{\pi}; I_{A\alpha}). \end{aligned}$$

Then  $\Gamma_{A\alpha}$  is well defined, and  $\Gamma_{A\alpha} = (\iota/\pi) \Psi_2$  in the notation of Theorem 2.5.

*Proof.* That  $(\iota/\pi) \Psi_2$  equals the above composite follows from the construction of  $\Psi_2$  in the proof of Theorem 2.5. That  $\Gamma_{A\alpha}$  is well defined thus follows automatically (but can also be shown directly, using Theorem 1.1).  $\blacksquare$

## 3

We still let  $A$  be a fixed unramified  $p$ -ring, with  $\varphi \in \text{Gal}(A/\hat{\mathbb{Z}}_p)$  and  $\text{Tr} = \text{Tr}_{A/\mathbb{Z}_p}$  as in Section 2. Theorem 2.5 will now be applied to calculate

$$K_2^*(A\pi) \cong \varprojlim \text{Ker}[K_1(A\tilde{\pi}, I_{A\alpha}) \rightarrow K_1(A\tilde{\pi})],$$

where  $\pi$  is a  $p$ -group and the limit is taken over all extensions

$$1 \rightarrow \rho \rightarrow \tilde{\pi} \xrightarrow{\alpha} \pi \rightarrow 1$$

of  $\pi$  by  $p$ -groups. For any such extension (not necessarily of  $p$ -groups), we set

$$I_{A\alpha} = \text{Ker}[A\tilde{\pi} \rightarrow A\pi] \quad \text{and} \quad M_{A\alpha} = \text{Ker}[\omega_{A\alpha}: H_0(\rho; I_{A\alpha}) \rightarrow \hat{\mathbb{Z}}_p \otimes \rho^{ab}]$$

(where  $\omega_{A\alpha}(\lambda(h-1)g) = \text{Tr}(\lambda) \otimes h$  for  $\lambda \in A$ ,  $h \in \rho$ , and  $g \in \tilde{\pi}$ ). Let

$$\iota_{A\alpha}: M_{A\alpha} \rightarrow H_0(\rho; I_{A\alpha}) \quad \text{and} \quad \kappa_{A\alpha}: H_0(\pi; M_{A\alpha}) \rightarrow \overline{I(A\tilde{\pi})}$$

be the maps induced by the inclusions. The main problem now is to study  $\text{Ker}(\kappa_{A\alpha})$ , and its limit over all such  $(\tilde{\pi}, \alpha)$ .

For simplicity, the subscripts in  $\kappa_{A\alpha}$ ,  $\omega_{A\alpha}$ , etc., will be dropped whenever possible.

**LEMMA 3.1.** *For any extension  $1 \rightarrow \rho \rightarrow \tilde{\pi} \xrightarrow{\alpha} \pi \rightarrow 1$  of groups there is an exact sequence*

$$\begin{aligned} H_1(\tilde{\pi}; I_{A\alpha}) &\xrightarrow{\Omega} H_1(\pi; \hat{\mathbb{Z}}_p \otimes \rho^{ab}) \rightarrow \text{Ker}[\kappa: H_0(\pi; M_{A\alpha}) \rightarrow \overline{I(A\tilde{\pi})}] \\ &\xrightarrow{\iota/\pi} \text{Ker}[H_0(\pi; I_{A\alpha}) \rightarrow \overline{I(A\tilde{\pi})}] \\ &\xrightarrow{\omega} \hat{\mathbb{Z}}_p \otimes \text{Ker}[\rho/[\tilde{\pi}, \rho] \rightarrow \pi^{ab}], \end{aligned}$$

where  $\Omega = \Omega_{A\alpha}$  is induced by  $\omega = \omega_{A\alpha}$ .

*Proof.* This follows directly from the homology sequence of the short exact sequence of  $\pi$ -modules

$$0 \rightarrow M_{A\alpha} \xrightarrow{\iota} H_0(\rho; I_{A\alpha}) \xrightarrow{\omega} \hat{\mathbb{Z}}_p \otimes \rho^{ab} \rightarrow 0.$$

Note that  $H_1(\tilde{\pi}; I_{A\alpha}) \rightarrow H_1(\pi; H_0(\rho; I_{A\alpha}))$  is onto (look at the spectral sequence). ■

The next lemma is standard, but will be needed frequently for handling homology with coefficients in permutation modules.



LEMMA 3.2. *Let  $G$  be a group, and  $S$  a set upon which  $G$  acts. Let  $A(S)$  denote the free  $A$ -module with basis  $S$ , regarded as an  $AG$ -module. Let  $X \subseteq S$  be a set of  $G$ -orbit representatives, and define, for all  $x \in X$ ,*

$$G_x = \{g \in G: g(x) = x\}.$$

*Then, for all  $n$ ,*

$$H_n(G; A(S)) \cong \sum_{x \in X} H_n(G_x; A).$$

*In particular,  $H_1(G; A(S))$  is generated by elements of the form  $g \otimes x$  for  $x \in S$  and  $g \in G_x$ .*

*Proof.* Just note that

$$A(S) \cong \sum_{x \in X} \text{Ind}_{G_x}^G(A),$$

so the result follows by Shapiro's lemma (see, e.g., [16, p. 131]). ■

Now let  $\pi$  be a  $p$ -group again. For any unramified  $p$ -ring  $A$ , we define a group  $\mathcal{U}(A\pi)$  as in the Introduction:

$$\mathcal{U}(A\pi) = H_1(\pi; A\pi) / \langle g \otimes \lambda g^n: g \in \pi, \lambda \in A, n \in \mathbb{Z} \rangle,$$

where  $\pi$  acts on  $A\pi$  via conjugation. By Lemma 3.2,

$$\mathcal{U}(A\pi) \cong \sum_i (Z_\pi(g_i) / S_\pi(g_i))^{ab} \otimes A(g_i),$$

where  $g_i$  runs over a set of conjugacy class representatives for  $\pi$  and

$$S_\pi(g) = \langle \sigma \in \pi: \sigma \text{ cyclic, } g \in \sigma \rangle = \langle h: g = h^n, \text{ some } n \rangle.$$

We write  $\mathcal{U}(\pi) = \mathcal{U}(\mathbb{Z}_p\pi)$  for short. Note that  $\mathcal{U}(A\pi) \cong A \otimes \mathcal{U}(\pi)$  in general.

PROPOSITION 3.3. *Let  $1 \rightarrow \rho \rightarrow \tilde{\pi} \xrightarrow{\alpha} \pi \rightarrow 1$  be any extension of  $p$ -groups such that  $\alpha(\sigma)$  is cyclic for any abelian  $\sigma \subseteq \pi$ . Define*

$$\theta_0: \mathcal{U}(A\pi) \rightarrow H_0(\tilde{\pi}; I_{A\alpha})$$

*by setting, for any  $\lambda \in A$ , commuting  $g, h \in \pi$ , and  $\tilde{g} \in \alpha^{-1}g, \tilde{h} \in \alpha^{-1}h$ ,*

$$\theta_0(h \otimes \lambda g) = \lambda(\tilde{h}\tilde{g}\tilde{h}^{-1} - \tilde{g}) \in I_{A\alpha} = \text{Ker}[A\tilde{\pi} \rightarrow A\pi].$$

*Then  $\theta_0$  is well defined, and induces an isomorphism*

$$\theta: \mathcal{U}(A\pi) \xrightarrow{\cong} \text{Ker}[H_0(\tilde{\pi}; I_{A\alpha}) \rightarrow \overline{H(A\tilde{\pi})}].$$

*Proof.* Let  $\tilde{\theta}$  denote the composite

$$\tilde{\theta}: H_1(\tilde{\pi}; A\pi) \rightarrow H_1(\pi; A(\pi)) \rightarrow \mathcal{U}(A\pi).$$

Consider the diagram

$$\begin{array}{ccccccc} H_1(\tilde{\pi}; A\tilde{\pi}) & \xrightarrow{\alpha_*} & H_1(\tilde{\pi}; A\pi) & \xrightarrow{\partial} & H_0(\tilde{\pi}; I_{A\alpha}) & \xrightarrow{i_*} & H_0(\tilde{\pi}; A\tilde{\pi}) \cong \overline{I(A\tilde{\pi})} \\ & & \searrow \tilde{\theta} & & \nearrow \theta_0 & & \\ & & & \mathcal{U}(A\pi) & & & \end{array}$$

where the row is induced by the short exact sequence

$$0 \rightarrow I_{A\alpha} \rightarrow A\tilde{\pi} \rightarrow A\pi \rightarrow 0$$

of  $\tilde{\pi}$ -modules. We defined  $\theta_0$  by the relation  $\theta_0(\tilde{\theta}(x)) = \partial(x)$ . To see that this is well defined and induces an isomorphism  $\mathcal{U}(A\pi) \cong \text{Ker}(i_*)$ , it will suffice to show that  $\text{Ker}(\tilde{\theta}) = \text{Im}(\alpha_*)$ .

By Lemma 3.2,

$$\text{Im}(\alpha_*) = \langle h \otimes \lambda \alpha(g) \in H_1(\tilde{\pi}; A\pi): \lambda \in A, h, g \in \tilde{\pi}, hg = gh \rangle.$$

On the other hand

$$\text{Ker}(\tilde{\theta}) = \langle h \otimes \lambda g \in H_1(\tilde{\pi}; A\pi): \lambda \in A, h \in \tilde{\pi}, g \in \pi, \text{ and } \alpha(h) \in S_\pi(g) \rangle.$$

So we must show that the subgroups

$$H_g = \langle h \in \tilde{\pi}: g \in \alpha Z_{\tilde{\pi}}(h) \rangle \subseteq \tilde{\pi} \quad \text{and} \quad \alpha^{-1} S_\pi(g) \subseteq \tilde{\pi}$$

are equal for all  $g \in \pi$ .

Clearly,  $\alpha^{-1} g \in H_g$ , and so  $\rho \subseteq \langle \alpha^{-1} g \rangle \subseteq H_g$ . It thus suffices to show that  $\alpha(H_g) = S_\pi(g)$ . But

$$\begin{aligned} \alpha(H_g) &= \langle h \in \pi: \tilde{h}\tilde{g} = \tilde{g}\tilde{h} \text{ for some } \tilde{h} \in \alpha^{-1}h, \tilde{g} \in \alpha^{-1}g \rangle \\ &= \langle h \in \pi: g, h \text{ generate a cyclic subgroup} \rangle \\ &= \langle h \in \pi: h^n = g, \text{ some } n \rangle = S_\pi(g). \quad \blacksquare \end{aligned}$$

In order to compute limits, we will need to know the existence of certain "sufficiently large" extensions of a given group; in particular, the existence of extensions meeting the hypotheses of Proposition 3.3. This is done in the following lemma:

LEMMA 3.4. For any  $p$ -group  $\pi$ , there is an extension

$$1 \rightarrow \rho \rightarrow \tilde{\pi} \xrightarrow{\alpha} \pi \rightarrow 1 \quad (1)$$

of  $p$ -groups having the properties

- (i) For any abelian  $\sigma \subseteq \tilde{\pi}$ ,  $\alpha(\sigma)$  is cyclic.
- (ii) For all  $n \geq 2$ , the boundary map

$$\partial_n: H_n(\pi) \rightarrow H_{n-2}(\tilde{\pi}; \rho^{ab})$$

(in the spectral sequence for (1)) is injective (note that  $\partial_2 = \delta^\alpha$ ).

In fact, if  $\pi = F/R$  where  $F$  is free, then  $\tilde{\pi}$  can be chosen of the form  $F/S$  for some  $S \subseteq R$ .

*Proof.* Write  $\pi = F/R$  where  $F$  is free and finitely generated. Let  $q = |\pi|$ , and set

$$S = \langle [R, R], R^q \rangle \quad \text{and} \quad \tilde{\pi} = F/S.$$

Then  $R/S \cong R^{ab} \otimes \mathbb{Z}/q$ , so  $\tilde{\pi}$  is a  $p$ -group. Let  $\alpha: \tilde{\pi} \rightarrow \pi$  be the projection.

(i) We claim that  $\alpha(\sigma)$  is cyclic for any abelian  $\sigma \subseteq \tilde{\pi}$ . To see this, choose  $g, h \in F$ , set  $F_0 = \langle R, g, h \rangle$ , and assume that  $F_0/R \subseteq \pi$  is not cyclic. We must show that  $[g, h] \notin S$ .

Since  $F_0/R$  is a non-cyclic  $p$ -group, we may choose  $R_0 \triangleleft F_0$  such that  $R_0 \cong R$ ,  $F_0/R_0 \cong \mathbb{Z}/p \times \mathbb{Z}/p$ , and  $F_0/R_0$  is generated by  $g$  and  $h$ . Then  $g \wedge h$  is non-zero in  $H_2(F_0/R_0) \cong \mathbb{Z}/p$ , and the composite

$$\begin{aligned} H_2(F_0/R_0) &\xrightarrow{\delta} \frac{R_0}{[F_0, R_0]} \rightarrow \frac{R_0}{[F_0, R_0]} \otimes \mathbb{Z}/p \\ &= R_0 / \langle [F_0, R_0], R_0^p \rangle \end{aligned}$$

is injective by Proposition 2.1(ii). Since  $\delta(g \wedge h) = [g, h]$  (by definition of  $g \wedge h$ ), this implies that

$$[g, h] \notin \langle [F_0, R_0], R_0^p \rangle \supseteq \langle [R, R], R^q \rangle = S.$$

(ii) Recall that  $\exp(H_2(\pi))$  divides  $|\pi| = q$  (see Proposition XII.2.5 in [3]). Hence

$$\delta^\alpha: H_2(\pi) \rightarrow H_0(\pi; (R/S)^{ab}) \cong (R/[R, F]) \otimes \mathbb{Z}/q$$

is injective by Proposition 2.1(ii). For  $n \geq 3$ ,

$$\partial: H_n(\pi) \rightarrow H_{n-2}(\pi; R^{ab})$$

is an isomorphism since  $H_i(F) \cong H_i(R) = 0$  for  $i \geq 2$ . But

$$\exp H_{n-2}(\pi; R^{ab}) \mid q$$

(see [3] again), so  $H_{n-2}(\pi; R^{ab})$  injects into  $H_{n-2}(\pi; R^{ab} \otimes \mathbb{Z}/q)$ , and we are done. ■

Lemma 3.4 can now be applied to describe some of the limits which arise in computing  $K_2^*(A\pi)$ .

**PROPOSITION 3.5.** *Let  $1 \rightarrow R \rightarrow F \rightarrow^\alpha \pi \rightarrow 1$  be an extension, where  $\pi$  is a  $p$ -group and  $F$  is free. Let  $\mathcal{S}$  be the set of all subgroups  $S \triangleleft F$  such that  $S \subseteq R$  and  $F/S$  is a  $p$ -group. For  $S \in \mathcal{S}$ , write  $\tilde{\pi}_S = F/S$ ,  $\rho_S = R/S$ , and let  $\alpha_S: \tilde{\pi}_S \rightarrow \pi$  be the projection. Then*

- (i)  $\varprojlim_{S \in \mathcal{S}} SK_1(A\tilde{\pi}_S) \cong \varprojlim_{S \in \mathcal{S}} SK_1(A\rho_S) = 0$ ,
- (ii)  $\varprojlim_{S \in \mathcal{S}} H_1(\pi; \rho_S^{ab}) \cong H_1(\pi; R^{ab}) \cong H_3(\pi)$ , and
- (iii)  $\varprojlim_{S \in \mathcal{S}} \text{Ker}[\kappa_{A\alpha_S}: H_0(\pi; M_{A\alpha_S}) \rightarrow \overline{I(A\tilde{\pi}_S)}] \\ \cong \text{Ker}[\kappa_{A\alpha}: H_0(\pi; M_{A\alpha}) \rightarrow \overline{I(AF)}].$

*Proof.* (i) Fix some  $S \in \mathcal{S}$ , set  $q = |\tilde{\pi}_S|$ , and define

$$S' = \langle [F, S], S^q \rangle \quad \text{and} \quad S'' = \langle [R, S], S^q \rangle.$$

Then  $S', S'' \in \mathcal{S}$ , since

$$S/S' \cong S/[F, S] \otimes \mathbb{Z}/q \quad \text{and} \quad S/S'' \cong S/[R, S] \otimes \mathbb{Z}/q$$

are  $p$ -groups. The boundary maps for the central extensions

$$1 \rightarrow S/S' \rightarrow \tilde{\pi}_{S'} \rightarrow \tilde{\pi}_S \rightarrow 1 \quad \text{and} \quad 1 \rightarrow S/S'' \rightarrow \rho_{S''} \rightarrow \rho_S \rightarrow 1$$

are injective by Proposition 2.1(ii), so  $SK_1(A\tilde{\pi}_{S'}) \cong SK_1(A\rho_{S''}) = 0$  by Proposition 2.1(iv), and the limits vanish.

(ii) Since  $H_n(F) \cong H_n(R) = 0$  for  $n \geq 2$ , the boundary map

$$\partial: H_3(\pi) \rightarrow H_1(\pi; R^{ab})$$

in the spectral sequence is an isomorphism. Furthermore, by Lemma 3.4,

$$\delta: H_2(\pi) \rightarrow H_0(\pi; \rho_S^{ab}) \quad \text{and} \quad \partial: H_3(\pi) \rightarrow H_1(\pi; \rho_S^{ab})$$

are injective for sufficiently small  $S \in \mathcal{S}$ . For such  $S$ , the spectral sequence for  $\rho_S \triangleleft \tilde{\pi}_S$  induces a surjection

$$H_2(\tilde{\pi}_S) \twoheadrightarrow \text{Coker}[\partial: H_3(\pi) \rightarrow H_1(\pi; \rho_S^{ab})],$$

and hence a surjection (all groups involved being finite)

$$0 = \varprojlim_{S \in \mathcal{S}} H_2(\tilde{\pi}_S) \rightarrow \text{Coker}[\varprojlim \partial: H_3(\pi) \rightarrow \varprojlim_{S \in \mathcal{S}} H_1(\pi; \rho_S^{ab})].$$

So  $\varprojlim(\partial)$  is an isomorphism.

(iii) Fix some  $S_0 \in \mathcal{S}$  such that  $\alpha_{S_0}: \tilde{\pi}_{S_0} \rightarrow \pi$  fulfills the hypotheses of Lemma 3.4. Then, for any  $S \subseteq S_0$  in  $\mathcal{S}$ ,

$$\text{Ker}[H_0(\tilde{\pi}_S; I_{A\alpha_S}) \rightarrow \overline{I(A\tilde{\pi}_S)}] \cong \text{Ker}[H_0(F; I_{A\alpha}) \rightarrow \overline{I(AF)}] \cong \mathcal{U}(A\pi)$$

by Proposition 3.3. Lemma 3.1 thus applies to give the following diagram with exact rows:

$$\begin{array}{ccccccc} H_1(F; I_{A\alpha}) & \longrightarrow & H_1(\pi; R^{ab}) & \longrightarrow & \text{Ker}(\kappa_{A\alpha}) & \longrightarrow & \mathcal{U}(A\pi) \xrightarrow{(\delta^\alpha)^{-1}\omega} H_2(\pi) \\ \downarrow f_S & & \downarrow & & \downarrow f'_S & & \downarrow \text{id} \\ H_1(F; I_{A\alpha_S}) & \longrightarrow & H_1(\pi; \rho_S^{ab}) & \longrightarrow & \text{Ker}(\kappa_{A\alpha_S}) & \longrightarrow & \mathcal{U}(A\pi) \longrightarrow H_2(\pi) \end{array}$$

Together with (ii), this implies that

$$f': \text{Ker}(\kappa_{A\alpha}) \rightarrow \varprojlim_{S \in \mathcal{S}} \text{Ker}(\kappa_{A\alpha_S})$$

is surjective. Furthermore, the diagram allows us to identify  $\text{Ker}(f'_S)$  with a subquotient of  $\text{Coker}(f_S)$ , so  $f'$  is an isomorphism if we can show that for every  $S \subseteq S_0$  in  $\mathcal{S}$ , there is  $S' \subseteq S$  in  $\mathcal{S}$  such that the map  $\text{Coker}(f_{S'}) \rightarrow \text{Coker}(f_S)$  is zero.

Fix  $S$ , and choose  $S' \subseteq S$  such that any abelian  $\sigma \subseteq \tilde{\pi}_S$  has cyclic image in  $\tilde{\pi}_{S'}$ . In particular, the induced map  $\mathcal{U}(\tilde{\pi}_{S'}) \rightarrow \mathcal{U}(\tilde{\pi}_S)$  is zero. The extensions

$$0 \rightarrow I_1 \rightarrow I_{A\alpha} \rightarrow I_{A\alpha_{S'}} \rightarrow 0 \quad \text{and} \quad 0 \rightarrow I_2 \rightarrow I_{A\alpha} \rightarrow I_{A\alpha_S} \rightarrow 0$$

(where  $I_1 = \text{Ker}[AF \rightarrow A\tilde{\pi}_{S'}]$ ,  $I_2 = \text{Ker}[AF \rightarrow A\tilde{\pi}_S]$ ) induce inclusions

$$\begin{aligned} \text{Coker}(f_{S'}) &\cong \text{Ker}[H_0(F; I_1) \rightarrow H_0(F; I_{A\alpha})] \subseteq \text{Ker}[H_0(F; I_1) \rightarrow \overline{I(AF)}] \\ &\cong \mathcal{U}(A\tilde{\pi}_{S'}) \end{aligned}$$

(applying Proposition 3.3), and similarly for  $\text{Coker}(f_S)$ . So  $\text{Coker}(f_{S'})$  maps trivially to  $\text{Coker}(f_S)$ . ■

These results can now be combined to describe  $K_2^*(A\pi)$  and  $Wh_2^*(A\pi)$ . Unfortunately, there does not seem to be any single formula or exact sequence providing a “best” description of these groups. So instead we give several descriptions.

For a  $p$ -group  $\pi$  and an unramified  $p$ -ring  $A$ , define

$$v = v_{A\pi}: \mathcal{U}(A\pi) \rightarrow H_2(\pi) \quad \text{and} \quad \tilde{v} = \tilde{v}_{A\pi}: \mathcal{U}(A\pi) \rightarrow A \otimes H_2(\pi)$$

by setting, for commuting  $g, h \in \pi$  and  $\lambda \in A$ ,

$$v(h \otimes \lambda g) = \text{Tr}(\lambda) \cdot (h \wedge g), \quad \tilde{v}(h \otimes \lambda g) = \lambda \otimes (h \wedge g)$$

(see Proposition 2.1(iii)). Set

$$\mathcal{U}_0(A\pi) = \text{Ker}[v: \mathcal{U}(A\pi) \rightarrow H_2(\pi)], \quad \mathcal{U}_0(\pi) = \mathcal{U}_0(\hat{\mathbb{Z}}_p \pi).$$

If  $\pi = F/R$  and  $\alpha: F \rightarrow \pi$  is the projection, then

$$I_{A\alpha} = \text{Ker}[AF \rightarrow A\pi] \quad \text{and} \quad M_{A\alpha} = \text{Ker}[\omega: H_0(R; I_{A\alpha}) \rightarrow \hat{\mathbb{Z}}_p \otimes R^{ab}]$$

as before. Let

$$\theta: \mathcal{U}(A\pi) \xrightarrow{\cong} \text{Ker}[H_0(F; I_{A\alpha}) \rightarrow \overline{I(AF)}]$$

again denote the isomorphism of Proposition 3.3.

**THEOREM 3.6.** *For any  $p$ -group  $\pi$ , there are exact sequences*

$$\begin{aligned} H_3(\pi) &\xrightarrow{\eta_{A\pi}} K_2^*(A\pi) \xrightarrow{(\omega_2, \Gamma_2^*)} (A \otimes H_2(\pi)) \oplus \mathcal{U}(A\pi) \\ &\xrightarrow[-\tilde{v}]{(1-\varphi) \otimes 1} A \otimes H_2(\pi), \end{aligned} \quad (1)$$

$$H_3(\pi) \xrightarrow{\eta_{A\pi}} Wh_2^*(A\pi) \xrightarrow{\Gamma_2^*} \mathcal{U}_0(A\pi) \rightarrow 0, \quad (2)$$

$$0 \rightarrow H_2(\pi) \rightarrow K_2^*(A\pi) \rightarrow Wh_2^*(A\pi) \rightarrow 0. \quad (3)$$

When  $A = \hat{\mathbb{Z}}_p$ , sequence (3) is split by  $\omega_2$ .

More precisely, if  $\pi = F/R$  and  $F$  is free and finitely generated, and  $\alpha: F \rightarrow \pi$  denotes the projection, there is an exact sequence

$$\begin{aligned} 0 \rightarrow K_2^*(A\pi) &\xrightarrow{(\omega_2, \Gamma_2^*)} (A \otimes H_2(\pi)) \oplus \text{Ker}[\kappa: H_0(\pi; M_{A\alpha}) \rightarrow \overline{I(AF)}] \\ &\xrightarrow{(1-\varphi) \otimes 1 - \xi} A \otimes H_2(\pi), \end{aligned} \quad (4)$$

where  $\xi$  is the composite

$$\begin{aligned} \text{Ker}(\kappa) &\xrightarrow{i/\pi} \text{Ker}[H_0(F; I_{A\alpha}) \rightarrow \overline{I(AF)}] \xrightarrow[\cong]{\theta^{-1}} \mathcal{U}(A\pi) \\ &\xrightarrow{\tilde{v}} A \otimes H_2(\pi) \end{aligned}$$

and  $\theta^{-1} \circ \iota / \pi \circ \tilde{I}_2^* = I_2^*$ . Also,  $\tilde{I}_2^*$  reduces to an isomorphism

$$Wh_2^*(A\pi) \xrightarrow{\cong} \text{Ker}[\kappa: H_0(\pi; M_{A\alpha}) \rightarrow \overline{I(AF)}]. \quad (5)$$

Furthermore, in (1) and (2),

$$\text{Ker}(\eta_{A\pi}) = \text{Im}[\Omega: H_1(F; I_\alpha) \rightarrow H_1(\pi; R^{ab}) \cong H_3(\pi)], \quad (6)$$

where  $I_\alpha = \text{Ker}[\mathbb{Z}F \rightarrow \mathbb{Z}\pi]$  and  $\Omega$  is induced by the map  $\omega = \omega_{\mathbb{Z}\alpha}: I_\alpha \rightarrow R^{ab}$ . In particular,  $\text{Ker}(\eta_{A\pi})$  is independent of  $A$ .

*Proof.* Let  $\mathcal{S}$ , and  $\rho_S, \tilde{\pi}_S, \alpha_S$  for  $S \in \mathcal{S}$ , be as in Proposition 3.5. The maps  $\alpha_S: \tilde{\pi}_S \rightarrow \pi$  are clearly cofinal among all surjections of  $p$ -groups onto  $\pi$ . It follows that

$$\begin{aligned} K_2^*(A\pi) &\cong \varprojlim_{S \in \mathcal{S}} \text{Coker}[K_2(A\tilde{\pi}_S) \rightarrow K_2(A\pi)] \\ &\cong \varprojlim_{S \in \mathcal{S}} \text{Ker}[K_1(A\tilde{\pi}_S, I_S) \rightarrow K_1(A\tilde{\pi}_S)] \quad (I_S = I_{A\alpha_S}) \\ &\cong \varprojlim_{S \in \mathcal{S}} \text{Ker}\left[\frac{K_1(A\tilde{\pi}_S, I_S)}{SK_1(A\rho_S)} \rightarrow \frac{K_1(A\tilde{\pi}_S; I(A\tilde{\pi}_S))}{SK_1(A\tilde{\pi}_S)}\right], \end{aligned}$$

where the last step holds by Proposition 3.5(i).

Hence, by Theorem 2.5 applied to the pairs  $\rho_S \triangleleft \tilde{\pi}_S$  and  $\tilde{\pi}_S \triangleleft \pi$ , there is an exact sequence

$$\begin{aligned} 0 \rightarrow K_2^*(A\pi) &\rightarrow A \otimes \varprojlim_{S \in \mathcal{S}} \text{Ker}[\rho_S / [\rho_S, \tilde{\pi}_S] \rightarrow \tilde{\pi}_S^{ab}] \oplus \varprojlim_{S \in \mathcal{S}} \text{Ker}(\kappa_{A\alpha_S}) \\ &\rightarrow A \otimes \varprojlim_{S \in \mathcal{S}} \text{Ker}[\rho_S / [\rho_S, \tilde{\pi}_S] \rightarrow \tilde{\pi}_S^{ab}]. \end{aligned} \quad (7)$$

But  $\varprojlim \text{Ker}(\kappa_{A\alpha_S}) = \text{Ker}(\kappa_{A\alpha})$  by Proposition 3.5, and

$$\varprojlim \text{Ker}[\rho_S / [\rho_S, \tilde{\pi}_S] \rightarrow \tilde{\pi}_S^{ab}] \cong \varprojlim \text{Coker}[H_2(\tilde{\pi}_S) \rightarrow H_2(\pi)] \cong H_2(\pi).$$

Substituting these into (7) gives sequence (4), where

$$\xi = (\delta^\alpha)^{-1} (\omega^* \iota) / \pi: \text{Ker}[\kappa: H_0(\pi; M_{A\alpha}) \rightarrow \overline{I(AF)}] \rightarrow A \otimes H_2(\pi).$$

To see that this equals the composite claimed in the Theorem, we check that the sequence

$$\begin{array}{ccc} \text{Ker}[H_0(F; I_{A\alpha}) \rightarrow \overline{I(AF)}] & \xrightarrow[\cong]{\theta^{-1}} & \mathcal{U}(A\pi) \\ \downarrow \omega^* / \pi & & \downarrow \bar{\gamma} \\ A \otimes \text{Ker}[R/[R, F] \rightarrow F^{ab}] & \xrightarrow[\cong]{(\delta^\alpha)^{-1}} & A \otimes H_2(\pi) \end{array} \quad (8)$$

commutes. But for any  $\lambda \in A$ , commuting  $g, h \in \pi$ , and  $\tilde{g} \in \alpha^{-1}g, \tilde{h} \in \alpha^{-1}h$ ,

$$\omega^* \theta(g \otimes \lambda h) = \omega^*(\lambda(\tilde{g}\tilde{h}\tilde{g}^{-1} - \tilde{h})) = \lambda \otimes [\tilde{g}, \tilde{h}] = \delta^\alpha(\tilde{v}(g \otimes \lambda h)).$$

Now consider the exact sequence of Lemma 3.1:

$$\begin{aligned} H_1(F; I_{A\alpha}) &\xrightarrow{\Omega} H_1(\pi; R^{ab}) \rightarrow \text{Ker}(\kappa_{A\alpha}) \xrightarrow{i/\pi} \text{Ker}[H_0(F; I_{A\alpha}) \rightarrow \overline{I(AF)}] \\ &\xrightarrow{(\delta^\alpha)^{-1}\omega} H_2(\pi). \end{aligned}$$

By the commutativity of (8), the composite

$$\mathcal{U}(A\pi) \xrightarrow{\theta} \text{Ker}[H_0(F; I_{A\alpha}) \rightarrow \overline{I(AF)}] \xrightarrow{(\delta^\alpha)^{-1}\omega} H_2(\pi)$$

is equal to  $v$  (recall that  $\omega = \text{Tr} \circ \omega^*$ ). Furthermore, the image of  $H_1(\pi; R^{ab})$  in  $\text{Ker}(\kappa_{A\alpha})$  is contained in  $\text{Ker}(\xi)$ , and hence in  $\text{Im}(I_2^*)$  in (4) above. So putting this into (4), we get a new exact sequence

$$\begin{aligned} H_1(\pi; R^{ab}) &\xrightarrow{\eta_{A\pi}} K_2^*(A\pi) \xrightarrow{(\omega_2, I_2^*)} (A \otimes H_2(\pi)) \oplus \mathcal{U}_0(A\pi) \\ &\xrightarrow[\sim]{(1-\varphi) \otimes 1} A \otimes H_2(\pi), \end{aligned}$$

where  $I_2^* = \theta^{-1}(i/\pi) \tilde{I}_2^*$ , and  $\text{Ker}(\eta_{A\pi}) = \text{Im}(\Omega)$ . Sequence (1) now follows upon identifying  $H_3(\pi)$  with  $H_1(\pi; R^{ab})$  and noting that  $(1-\varphi)A \subseteq \text{Ker}(\text{Tr})$  (so that replacing  $\mathcal{U}_0(A\pi)$  by  $\mathcal{U}(A\pi)$  does not change exactness).

By definition (and the fact that  $K_2(A)$  is generated by symbols [13]),

$$Wh_2(A\pi) = \text{Coker}[\lambda_\pi: H_2(\pi) \rightarrow K_2(A\pi)/\{A^*, A^* \times \pi^{ab}\}].$$

Hence  $Wh_2^*(A\pi) \cong K_2^*(A\pi)/\text{Im}(\lambda_\pi)$ : the symbols  $\{A^*, A^* \times \pi^{ab}\}$  clearly vanish in  $K_2^*(A\pi)$ . By Theorem 2.5 (and the construction of sequence (4)), the composite

$$H_2(\pi) \xrightarrow{\lambda_\pi} K_2^*(A\pi) \xrightarrow{(\omega_2, I_2^*)} (A \otimes H_2(\pi)) \oplus \text{Ker}(\kappa_{A\alpha})$$

is inclusion into the first factor. So (3) is exact ( $H_2(\pi)$  injects into  $K_2^*(A\pi)$ ), (2) now follows easily from (1), and (5) from (4).

We have seen that  $\text{Ker}(\eta_{A\pi}) = \text{Im}(\Omega_{A\alpha})$  where  $\Omega_{A\alpha}$  by definition is the composite

$$\begin{aligned} H_1(F; I_{A\alpha}) &\cong A \otimes_{\mathbb{Z}} H_1(F; I_\alpha) \xrightarrow{\text{Tr} \otimes 1} \hat{\mathbb{Z}}_p \otimes_{\mathbb{Z}} H_1(F; I_\alpha) \\ &\xrightarrow{\Omega_\alpha} H_1(\pi; R^{ab}) \cong H_3(\pi). \end{aligned}$$



Since the trace map is onto and  $H_3(\pi)$  is a  $p$ -group, this implies that  $\text{Im}(\Omega_{A\alpha}) = \text{Im}(\Omega_\alpha)$ , and in particular,  $\text{Im}(\Omega_{A\alpha})$  is independent of  $A$ . ■

For future reference, we now give explicit descriptions of the maps  $\omega_2$  and  $\Gamma_2^*$  occurring in Theorem 3.6.

PROPOSITION 3.7. *The maps*

$$\omega_2(A\pi): K_2^*(A\pi) \rightarrow A \otimes H_2(\pi) \quad \text{and} \quad \Gamma_2^*(A\pi): K_2^*(A\pi) \rightarrow \mathcal{U}(A\pi)$$

are described as follows:

(i) *Let  $1 \rightarrow \rho \rightarrow \tilde{\pi} \xrightarrow{\alpha} \pi \rightarrow 1$  be any extension of  $p$ -groups such that  $\delta^\alpha$  is injective. Then  $\omega_2(A\pi)$  is the composite*

$$\begin{aligned} K_2(A\pi) &\xrightarrow{\partial} \text{Ker}[K_1(A\tilde{\pi}, I_{A\alpha}) \rightarrow K_1(A\tilde{\pi})] \\ &\xrightarrow{\omega^*} A \otimes \text{Ker}[\rho/[\tilde{\pi}, \rho] \rightarrow \pi^{ab}] \xleftarrow{\delta^\alpha} A \otimes H_2(\pi). \end{aligned}$$

(ii) *Let  $1 \rightarrow \rho \rightarrow \tilde{\pi} \xrightarrow{\alpha} \pi \rightarrow 1$  be any extension of  $p$ -groups such that  $\alpha(\sigma)$  is cyclic for any abelian  $\sigma \subseteq \pi$ . Then  $\Gamma_2^*(A\pi)$  is the composite*

$$\begin{aligned} K_2(A\pi) &\xrightarrow{\partial} \text{Ker}[K_1(A\tilde{\pi}, I_{A\alpha}) \rightarrow K_1(A\tilde{\pi})] \\ &\xrightarrow{\Gamma_{A\alpha}} \text{Ker}[H_0(\tilde{\pi}, I_{A\alpha}) \rightarrow \overline{I(A\tilde{\pi})}] \xleftarrow{\theta} \mathcal{U}(A\pi), \end{aligned}$$

where  $\Gamma_{A\alpha}$  is the map defined in Proposition 2.6.

*Proof.* These follow directly from Theorem 2.5, Proposition 2.6, and the construction of sequence (1) in Theorem 3.6. ■

Finally, as a more concrete example, we show:

PROPOSITION 3.8. *Let  $\pi$  be a  $p$ -group and  $A$  an unramified  $p$ -ring. Then, for any  $g \in \pi$  and  $u \in 1 + I(A[Z_\pi(g)])$ ,*

$$\Gamma_2^*(\{g, u\}) = g \otimes \Gamma(u) \in \mathcal{U}(A\pi) \quad (\Gamma = \Gamma_{A[Z_\pi(g)]}).$$

*Proof.* Let  $Z = Z_\pi(g)$ . Fix an extension  $1 \rightarrow \rho \rightarrow \tilde{\pi} \xrightarrow{\alpha} \pi \rightarrow 1$  of  $p$ -groups such that  $\alpha(\sigma)$  is cyclic for all abelian  $\sigma \subseteq \tilde{\pi}$ . Lift  $g$  to  $\tilde{g} \in \tilde{\pi}$  and  $u$  to

$$\tilde{u} \in 1 + I(A[\alpha^{-1}Z]).$$

Then  $\partial(\{g, u\}) = [\tilde{g}, \tilde{u}] \in K_1(A\tilde{\pi}, I_{A\alpha})$ .

Let  $G \subseteq \tilde{\pi} \times \tilde{\pi}$  be the pullback over  $\pi$ , and define maps

$$\Delta: A\tilde{\pi} \rightarrow AG \quad \text{and} \quad \Delta_{\tilde{g}}: A[\alpha^{-1}Z] \rightarrow AG$$

by setting  $\Delta(h) = (h, h)$ ,  $\Delta_{\tilde{g}}(h) = (\tilde{g}\tilde{h}\tilde{g}^{-1}, \tilde{h})$ . Then  $[\tilde{g}, \tilde{u}]$  lifts to

$$\Delta_{\tilde{g}}(\tilde{u}) \Delta(\tilde{u})^{-1} \in \text{Ker}[r_2: K_1(AG) \rightarrow K_1(A\tilde{\pi})]$$

in the notation of Proposition 2.6, and so

$$\Gamma_{A\alpha}([\tilde{g}, \tilde{u}]) = r_1(\Delta_{\tilde{g}}(\Gamma(\tilde{u})) - \Delta(\Gamma(\tilde{u}))) = \tilde{g}\Gamma(\tilde{u})\tilde{g}^{-1} - \Gamma(\tilde{u}) \in H_0(\tilde{\pi}; I_{A\alpha}).$$

Hence  $\Gamma_2^*(\{g, u\}) = g \otimes \Gamma(u)$  by Proposition 3.7 (and the definition of  $\theta$  in Proposition 3.3). ■

#### 4

The “mysterious” part of  $Wh_2^*(A\pi)$ , under the description given in Theorem 3.6, is the group

$$\text{Im}[\eta_{A\pi}: H_3(\pi) \rightarrow Wh_2^*(A\pi)] = \text{Ker}[\Gamma_2^*: Wh_2^*(A\pi) \rightarrow \mathcal{U}_0(A\pi)].$$

This will now be studied; the main results of this section are that  $\eta_{A\pi} = 0$  for abelian  $\pi$ , but that  $\eta_{A\pi}$  is not zero in general. Recall (Theorem 3.6) that

$$\text{Ker}(\eta_{A\pi}) = \text{Im}[\Omega_{\pi}: H_1(F; I_{\alpha}) \rightarrow H_1(\pi; R^{ab}) \cong H_3(\pi)]$$

when  $\pi = F/R$ ,  $\alpha: F \rightarrow \pi$  is the projection, and  $F$  is free.

We start with a closer look at  $H_1(F; I_{\alpha})$ .

LEMMA 4.1. *Let  $\pi = F/R$  be a  $p$ -group, where  $F$  is free and  $\alpha: F \rightarrow \pi$  denotes the projection. Set*

$$I_{\alpha} = \text{Ker}[\alpha: \mathbb{Z}F \rightarrow \mathbb{Z}\pi].$$

Then

$$H_1(F; I_{\alpha}) \cong \text{Ker}[H_1(F; \mathbb{Z}F) \rightarrow H_1(F; \mathbb{Z}\pi)],$$

and is generated by elements of the form  $\sum_{i=1}^k a_i \otimes b_i \in H_1(F; \mathbb{Z}F)$  where  $a_i, b_i \in F$ , and

- (i)  $\alpha(b_i) = g$  for some fixed  $g \in \pi$  and all  $i$ ,
- (ii)  $\langle a_i, b_i \rangle \subseteq F$  is cyclic for all  $i$ , and
- (iii)  $a_1 a_2 \cdots a_k = 1$  in  $(\alpha^{-1}Z_{\pi}(g))^{\text{ab}}$ .

*Proof.* Let  $g_1, \dots, g_n$  be conjugacy class representatives for  $\pi$ . By Lemma 3.2,

$$H_2(F; \mathbb{Z}\pi) \cong \sum_i H_2(\alpha^{-1}Z_\pi(g_i)) = 0$$

( $\alpha^{-1}Z_\pi(g_i) \subseteq F$  is free for all  $i$ ), and so

$$H_1(F; I_\alpha) \cong \text{Ker}[\alpha_*: H_1(F; \mathbb{Z}F) \rightarrow H_1(F; \mathbb{Z}\pi)]$$

by the homology exact sequence.

Applying Lemma 3.2 again,  $\text{Ker}(\alpha_*)$  splits as a sum

$$\sum_{j=1}^n \text{Ker}[H_1(\alpha^{-1}Z_\pi(g_j); \mathbb{Z}(\alpha^{-1}g_j)) \rightarrow H_1(\alpha^{-1}Z_\pi(g_j); \mathbb{Z})].$$

Furthermore, also by Lemma 3.2,  $H_1(\alpha^{-1}Z_\pi(g_j); \mathbb{Z}(\alpha^{-1}g_j))$  is generated by elements  $a \otimes b$  such that  $\alpha(b) = g_j$  and  $[a, b] = 1$ . In particular,  $\langle a, b \rangle$  is an abelian subgroup of  $F$ , and hence cyclic. So  $\text{Ker}(\alpha_*)$  is generated by elements  $\sum_{i=1}^k a_i \otimes b_i$ , such that  $\alpha(b_i) = g_j$  for all  $i$  and some fixed  $j$ , such that  $\langle a_i, b_i \rangle$  is cyclic for all  $i$ , and such that

$$\alpha_* \left( \sum_{i=1}^k a_i \otimes b_i \right) = \sum_{i=1}^k a_i \otimes g_j = \left( \prod_{i=1}^k a_i \right) \otimes g_j = 1 \quad (1)$$

in  $H_1(\alpha^{-1}Z_\pi(g_j); \mathbb{Z}(g_j))$ . But (1) holds if and only if

$$a_1 \cdots a_k = 1 \quad \text{in } (\alpha^{-1}Z_\pi(g_j))^{ab}. \quad \blacksquare$$

In order to make explicit homology calculations, we will use the bar construction as defined, for example, in Section VI.13 of [6], but reflected so as to get a resolution of free *right* modules. In other words, for a group  $G$  and left  $\mathbb{Z}G$ -module  $M$ , we let  $B'_n(G; M)$  be the sum of terms

$$[g_1 | g_2 | \cdots | g_n] \otimes M$$

for all  $(g_1, \dots, g_n) \in (G-1)^n$ . As usual, we set  $[g_1 | \cdots | g_n] \otimes M = 0$  if  $g_i = 1$  for some  $i$ . The boundary maps in lower dimensions are given by

$$\partial([g | h] \otimes x) = [g] \otimes hx - [gh] \otimes x + [h] \otimes x$$

and

$$\partial([g] \otimes x) = gx - x$$

for  $g, h \in G$  and  $x \in M$ . In particular, there is the relation

$$[gh] \otimes x \equiv [h] \otimes x + [g] \otimes hx \quad (\text{mod boundaries}) \quad (*)$$

in  $B'_1(G; M)$ . As before,  $[g] \otimes x$  will be written  $g \otimes x$  for short.

The main result of this section can now be shown:

**THEOREM 4.2.** *If  $\pi$  is an abelian  $p$ -group, then  $\eta_{A\pi} = 0$  for any  $A$ . Hence*

$$Wh_2^*(A\pi) \cong \mathcal{U}_0(A\pi) \quad \text{and} \quad K_2^*(\hat{\mathbb{Z}}_p \pi) \cong \mathcal{U}_0(\pi) \oplus H_2(\pi)$$

in this case.

*Proof.* Write  $\pi = \mathbb{Z}/p^{n_1} \times \cdots \times \mathbb{Z}/p^{n_t}$ , where  $n_1 \geq n_2 \geq \cdots \geq n_t$ , and let  $g_1, \dots, g_t$  be the corresponding generators. By the Künneth formula,

$$\begin{aligned} H_3(\pi) &\cong \sum_{i=1}^t H_3(\mathbb{Z}/p^{n_i}) \oplus \sum_{i < j < k} \mathbb{Z}/p^{n_i} \otimes \mathbb{Z}/p^{n_j} \otimes \mathbb{Z}/p^{n_k} \\ &\quad \oplus \sum_{i < j} \text{Tor}(\mathbb{Z}/p^{n_i}, \mathbb{Z}/p^{n_j}) \\ &\cong \pi \oplus A^2(\pi) \oplus A^3(\pi) \end{aligned}$$

(where the isomorphisms are, of course, non-canonical).

Let  $F$  be the free group on generators  $a_1, \dots, a_t$ , let  $\alpha: F \rightarrow \pi$  be the map defined by  $\alpha(a_i) = g_i$ , and set  $R = \text{Ker}(\alpha)$ . Let

$$\Omega: H_1(F; I_\alpha) \rightarrow H_1(\pi; R^{ab}) \cong H_3(\pi)$$

be the homomorphism of Theorem 3.6; we must show that  $\Omega$  is onto. To do this, subgroups  $L_1, L_2, L_3 \subseteq H_1(F; I_\alpha)$  are defined as follows:

$$L_1 = \langle x_i = a_i \otimes (a_i^q - 1): 1 \leq i \leq t, q = p^{n_i} = |g_i| \rangle,$$

$$L_2 = \langle y_{ij} = a_i a_j \otimes ((a_i a_j)^q - 1): 1 \leq i < j \leq t, q = p^{n_i} = |g_i g_j| \rangle,$$

and

$$L_3 = \langle z_{ijk} = [a_i, a_j] a_k \otimes [a_i, a_j] a_k - a_k \otimes a_k: 1 \leq i < j < k \leq t \rangle.$$

In defining  $L_3$  we are of course identifying

$$H_1(F; I_\alpha) \cong \text{Ker}[\alpha_*: H_1(F; \mathbb{Z}F) \rightarrow H_1(F; \mathbb{Z}\pi)].$$

Now let  $\beta: R^{ab} \rightarrow R/[F, R]$  be the projection, and

$$\beta_*: H_1(\pi; R^{ab}) \rightarrow H_1(\pi; R/[F, R]) \cong \pi \otimes R/[F, R]$$

the induced map. Then for all  $i$  and  $j$ ,

$$\beta_* \Omega(x_i) = g_i \otimes a_i^q \quad (q = p^{n_i})$$

and

$$\beta_* \Omega(y_{ij}) = g_i g_j \otimes (a_i a_j)^q \quad (q = p^{n_i}).$$

Furthermore, using relation (\*) above, we get (for  $1 \leq i < j < k \leq t$ )

$$\begin{aligned} z_{ijk} &= [a_i, a_j] a_k \otimes ([a_i, a_j] - 1) a_k + [a_i, a_j] \otimes a_k \\ &= [a_i, a_j] a_k \otimes ([a_i, a_j] - 1) a_k \\ &\quad + a_i \otimes (a_j a_i^{-1} a_j^{-1} a_k a_j a_i^{-1} - a_i^{-1} a_j^{-1} a_k a_j a_i) \\ &\quad + a_j \otimes (a_i^{-1} a_j^{-1} a_k a_j a_i - a_j^{-1} a_k a_j) \in H_1(F; I_\alpha), \end{aligned}$$

and hence

$$\begin{aligned} \beta_* \Omega(z_{ijk}) &= g_k \otimes [a_i, a_j] + g_i \otimes [a_j, a_i^{-1} a_j^{-1} a_k a_j a_i] + g_j \otimes [a_i^{-1}, a_j^{-1} a_k a_j] \\ &= g_k \otimes [a_i, a_j] + g_i \otimes [a_j, a_k] + g_j \otimes [a_k, a_i] \in \pi \otimes R/[F, R]. \end{aligned}$$

Since  $R/(R \cap [F, F]) \cong F^{ab}$  is free abelian,  $R/[F, R]$  splits as a sum

$$R/[F, R] \cong \frac{R \cap [F, F]}{[R, F]} \oplus \frac{R}{R \cap [F, F]} \cong H_2(\pi) \oplus \frac{R}{R \cap [F, F]}.$$

Comparison of the elements  $\beta_* \Omega(x_i)$ ,  $\beta_* \Omega(y_{ij})$ , and  $\beta^* \Omega(z_{ijk})$  shows that

$$\beta_* \Omega(L_1 + L_2 + L_3) \cong \pi \oplus \Lambda^2(\pi) \oplus \Lambda^3(\pi) \cong H_3(\pi),$$

and hence that  $\Omega$  is onto. ■

We have been unable to get any feel for how “complicated” a group  $\pi$  must be in order that  $\eta_{A\pi} \neq 0$ . So the examples constructed in the following proposition are probably far from minimal.

**PROPOSITION 4.3.** *For any prime  $p$  there is a  $p$ -group  $\pi$  such that  $\eta_{A\pi} \neq 0$ , i.e., such that  $|Wh_2^*(A\pi)| > |\mathcal{U}_0(A\pi)|$ , for all  $A$ .*

*Proof.* Fix a prime  $p$ , and set  $q = p$  if  $p$  is odd,  $q = 4$  if  $p = 2$ . Let  $F$  be the free group on three generators  $a, b, c$ , and define subgroups

$$R = \langle [F, F], a^q, b^q, c^q \rangle$$

and

$$S = \langle [F, R], a^{-q}[b, c], b^{-q}[c, a], c^{-q}[a, b] \rangle.$$

Set  $\pi = F/R \cong (\mathbb{Z}/q)^3$  and  $\tilde{\pi} = F/S$ ; there is a central extension

$$1 \rightarrow R/S \rightarrow \tilde{\pi} \xrightarrow{\beta} \pi \rightarrow 1$$

where  $R/S \cong (\mathbb{Z}/q)^3$ . We will show that  $\eta_{A\tilde{\pi}} \neq 0$ .

To do this, we define another subgroup

$$T = \langle [F, R], a^q, b^q, c^q, [a, b]^p, [a, c]^p, [b, c]^p \rangle,$$

set  $G = F/T$ , and consider the central extension

$$1 \rightarrow R/T \rightarrow G \xrightarrow{\gamma} \pi \rightarrow 1 \quad (R/T \cong (\mathbb{Z}/p)^3).$$

Note in particular:

(i) any cyclic subgroup of  $\pi$  lifts to  $G$ : for example,

$$(ab)^q = a^q b^q [a, b]^{q(q-1)/2} = 1 \quad \text{in } G \quad (p \mid \tfrac{1}{2}q(q-1));$$

(ii) if  $p = 2$ , then any subgroup of  $\pi$  ( $\cong (\mathbb{Z}/4)^3$ ) isomorphic to  $\mathbb{Z}/4 \times \mathbb{Z}/2 \times \mathbb{Z}/2$  lifts to  $G$ : any liftings of elements of orders two and four in  $\pi$  commute in  $G$ .

We now claim that

$$\beta_*(\text{Ker}(\eta_{A\bar{\pi}})) \subseteq \text{Im}[\gamma_*: H_3(G) \rightarrow H_3(\pi)]. \quad (1)$$

Let  $\alpha: F \rightarrow \pi$  and  $\tilde{\alpha}: F \rightarrow \tilde{\pi}$  be the projections, and identify

$$H_1(F; I_{\tilde{\alpha}}) \cong \text{Ker}[H_1(F; \mathbb{Z}F) \rightarrow H_1(F; \mathbb{Z}\tilde{\pi})]$$

as in Lemma 4.1. Fix  $g \in \tilde{\pi}$  and  $\sum_{i=1}^k a_i \otimes b_i \in H_1(F; I_{\tilde{\alpha}})$ , where  $a_i, b_i \in F$ ,  $\alpha(b_i) = g$ , and  $\langle a_i, b_i \rangle \subseteq F$  is cyclic for all  $i$ , and  $a_1 \cdots a_k = 1$  in  $(\alpha^{-1}Z)^{ab}$  ( $Z = Z_{\tilde{\pi}}(g)$ ). Then

$$\Omega_{\tilde{\pi}}\left(\sum a_i \otimes b_i\right) \in \text{Im}[H_3(Z) \rightarrow H_3(\tilde{\pi})],$$

and so

$$\beta_* \Omega_{\tilde{\pi}}\left(\sum a_i \otimes b_i\right) \in \text{Im}[H_3(\beta Z) \rightarrow H_3(\pi)].$$

If  $g \notin R/S$ , then  $\beta(Z) = \beta(Z_{\tilde{\pi}}(g))$  is either cyclic or isomorphic to  $\mathbb{Z}/4 \times (\mathbb{Z}/2)^2$ , and hence lifts to  $G$  by (i) or (ii) above. So in this case,

$$\beta_* \Omega_{\tilde{\pi}}\left(\sum a_i \otimes b_i\right) \in \text{Im}[H_3(G) \rightarrow H_3(\pi)].$$

Now assume  $g \in R/S$ . Then  $b_i \in R$  for all  $i$ , so

$$\sum a_i \otimes b_i = \sum a_i \otimes (b_i - 1) \in H_1(F; I_{\alpha}) \cong \text{Ker}[H_1(F; \mathbb{Z}F) \rightarrow H_1(F; \mathbb{Z}\pi)],$$

where now  $(b_i - 1) \in I_\alpha$  for all  $i$ . Then

$$\beta_* \Omega_{\bar{\pi}} \left( \sum a_i \otimes b_i \right) = \Omega_{\pi} \left( \sum a_i \otimes (b_i - 1) \right) = \sum a_i \otimes b_i \in H_1(\pi; R^{ab}) \cong H_3(\pi).$$

Since  $\langle a_i, b_i \rangle \subseteq F$  is cyclic for each  $i$ , each term  $a_i \otimes b_i$  is a homology element, and in fact lies in the image of  $H_3(\sigma_i)$  for some cyclic  $\sigma_i \subseteq \pi$ . By (i), each  $\sigma_i$  lifts to  $G$ , and so again

$$\beta_* \Omega_{\bar{\pi}} \left( \sum a_i \otimes b_i \right) \in \text{Im}[H_3(G) \rightarrow H_3(\pi)].$$

Hence (1) follows upon combining these two cases:

$$\beta_*(\text{Ker}(\eta_{A\bar{\pi}})) = \beta_* \Omega_{\bar{\pi}}(H_1(F; I_{\bar{\alpha}})) \subseteq \text{Im}[\gamma_*: H_3(G) \rightarrow H_3(\pi)].$$

Finally, consider the element

$$\begin{aligned} x = & bab^{-1} \otimes [b^{-1}, bcb^{-1}](bab^{-1})^q + cbc^{-1} \otimes [c^{-1}, cac^{-1}](cbc^{-1})^q \\ & + aca^{-1} \otimes [a^{-1}, aba^{-1}](aca^{-1})^q \in H_1(\bar{\pi}; S^{ab}) \cong H_3(\bar{\pi}). \end{aligned} \quad (2)$$

That  $x$  actually is a cycle follows from the Hall commutator identity

$${}^b[a, [b^{-1}, c]] \cdot {}^c[b, [c^{-1}, a]] \cdot {}^a[c, [a^{-1}, b]] = 1 \quad \text{in } F.$$

We claim that  $\beta_*(x) \notin \text{Im}(\gamma_*)$ , and hence by (1) that  $x \notin \text{Ker}(\eta_{A\bar{\pi}})$ . Identify

$$H_3(G) \cong H_1(G; T^{ab}) \quad \text{and} \quad H_3(\pi) \cong H_1(\pi; R^{ab});$$

then

$$\text{Im}[\gamma_*: H_1(G; T^{ab}) \rightarrow H_1(\pi; R^{ab})] \subseteq \text{Ker}[\partial: H_1(\pi; R^{ab}) \rightarrow \pi \otimes R/T].$$

Upon simplifying (2), we get

$$\partial(\beta_* x) = -a \otimes [b, c] - b \otimes [c, a] - c \otimes [a, b] \in \pi \otimes R/T$$

(recall that  $a^q, b^q, c^q \in T$ ). But  $\{[b, c], [c, a], [a, b]\}$  is a basis for  $R/T \cong (\mathbb{Z}/p)^3$ , so  $\partial(\beta_* x) \neq 0$  and  $\beta_*(x) \notin \text{Im}(\gamma_*)$ . ■

To finish the section, we show:

**PROPOSITION 4.4.** *For any prime  $p$ , unramified  $p$ -ring  $A$ , and  $p$ -group  $\pi$ ,  $K_2^*(A\pi) = 0$  if and only if  $\pi$  is cyclic or quaternionic.*

*Proof.* Assume first that  $\pi$  is not cyclic or quaternionic. Fix any  $b \in \mathbb{Z}(\pi)$  of order  $p$ . Using Theorem 5.4.10 in [5], choose any  $a \notin \langle b \rangle$  in  $\pi$ .

also of order  $p$ . Choose  $g \in \pi$  such that  $g^{p^k} = a$  for some  $k$ , but such that  $g$  is not a  $p$ -power in  $\pi$ . Then

$$S_\pi(g) = \langle \sigma \subseteq \pi \mid \sigma \text{ cyclic, } g \in \sigma \rangle = \langle g \rangle.$$

So  $b \in Z_\pi(g) - S_\pi(g)$ ,  $(Z_\pi(g)/S_\pi(g))^{ab} \neq 1$ , and so  $\mathcal{U}(\pi) \neq 1$ . But

$$\mathcal{U}_0(A\pi) = \text{Ker}[\text{Tr} \otimes v: A \otimes \mathcal{U}(\pi) \rightarrow H_2(\pi)],$$

and so by Theorem 3.6

$$|K_2^*(A\pi)| \geq |H_2(\pi)| \cdot |\mathcal{U}_0(A\pi)| \geq |A \otimes \mathcal{U}(\pi)| > 1.$$

If  $\pi$  is cyclic, then  $K_2^*(A\pi) = 0$  by Theorem 3.6 and 4.2 (or just from the fact that  $K_2(A\pi)$  is generated by symbols). If  $\pi$  is quaternionic, then  $H_2(\pi) = 0$  (see Section XII.7 of [3]), and one easily checks that  $\mathcal{U}(\pi) = 0$ . So by Theorem 3.6, it remains only to check that  $\eta_{A\pi} = 0$  in this case.

Set  $2^n = |\pi|$  ( $n \geq 3$ ), and write  $\pi = F/R$  where  $F = \langle a, b \rangle$  and  $R$  is the normal subgroup generated by

$$\{a^{2^{n-1}}, a^{2^{n-2}}b^{-2}, bab^{-1}a\}.$$

Let  $\alpha: F \rightarrow \pi$  be the projection. Consider the element

$$\begin{aligned} x &= a \otimes a^{2^{n-2}} + b \otimes b^2 - ab \otimes (ab)^2 \in \text{Ker}[H_1(F; \mathbb{Z}F) \rightarrow H_1(F; \mathbb{Z}\pi)] \\ &\cong H_1(F; I_\alpha) \end{aligned}$$

(see Lemma 4.1). Modulo boundaries,

$$\begin{aligned} x &= a \otimes a^{2^{n-2}} + b \otimes b^2 - b \otimes (ab)^2 - a \otimes b(ab)^2 b^{-1} \\ &= a \otimes (a^{2^{n-2}} - (ba)^2) + b \otimes (b^2 - (ab)^2) \end{aligned}$$

(see relation (\*) above), and so

$$\Omega(x) = a \otimes a^{2^{n-2}}(ba)^{-2} + b \otimes b^2(ab)^{-2} \in H_1(\pi; R^{ab}) \cong H_3(\pi).$$

Now let

$$\begin{aligned} \beta_*: H_1(\pi; R^{ab}) &\rightarrow H_1\left(\pi; \frac{R}{R \cap [F, F]}\right) \cong \pi^{ab} \otimes \frac{R}{R \cap [F, F]} \\ &\cong (\mathbb{Z}/2)^2 \otimes \langle a^2, b^2 \rangle \end{aligned}$$

be the projection. Then

$$\beta_* \Omega(x) = a \otimes a^{2^{n-2}-2}b^{-2} + b \otimes a^{-2}$$



is not a multiple of two in  $\pi^{ab} \otimes R/(R \cap [F, F])$ . So  $\Omega(x)$  must generate  $H_3(\pi) \cong \mathbb{Z}/2^n$  [3, Sect. XII.7],  $\Omega$  is onto, and hence  $\eta_{A\pi} = 0$  by Theorem 3.6(5). ■

## 5

These results will now be applied to get lower bounds on the size of  $K_2(\mathbb{Z}\pi)$  and  $Wh_2(\pi)$  for  $p$ -groups  $\pi$ . The main problem is to get an appropriate localization sequence for relating  $K_2(\mathbb{Z}\pi)$  to  $K_2^*(\widehat{\mathbb{Z}}_p\pi)$ .

**THEOREM 5.1.** *Let  $\mathfrak{A} \subseteq \mathfrak{B}$  be  $\mathbb{Z}$ -orders in a semisimple  $\mathbb{Q}$ -algebra  $A$ . Then, for any prime  $p$ , there is an exact sequence*

$$0 \rightarrow \varinjlim_n \text{Coker}[K_2(\mathfrak{A}) \rightarrow K_2(\mathfrak{A}/p^n\mathfrak{A})] \rightarrow \varinjlim_n \text{Coker}[K_2(\mathfrak{B}) \rightarrow K_2(\mathfrak{B}/p^n\mathfrak{B})] \\ \rightarrow Cl_1(\mathfrak{A})_{(p)} \rightarrow Cl_1(\mathfrak{B})_{(p)} \rightarrow 0.$$

*Proof.* This follows from Proposition 1.2 in [11], except for showing that the map

$$\varphi: \varinjlim_n \text{Coker}[K_2(\mathfrak{A}) \rightarrow K_2(\mathfrak{A}/p^n\mathfrak{A})] \rightarrow \varinjlim_n \text{Coker}[K_2(\mathfrak{B}) \rightarrow K_2(\mathfrak{B}/p^n\mathfrak{B})]$$

is injective.

*Step 1.* We first show that the map

$$\varphi_0: \varinjlim_I K_1(\mathfrak{A}, I) \rightarrow \varinjlim_I K_1(\mathfrak{B}, I)$$

is injective, where the limits are taken over all ideals of finite index. It suffices to show for any  $\mathfrak{B}$ -ideal  $I \subseteq \mathfrak{A}$  of finite index that there exists  $I' \subseteq I$  such that the map

$$\text{Ker}[K_1(\mathfrak{A}, I') \rightarrow K_1(\mathfrak{B}, I')] \rightarrow \text{Ker}[K_1(\mathfrak{A}, I) \rightarrow K_1(\mathfrak{B}, I)]$$

is zero. By stability results [2, Corollary I.4.5], this means finding  $I'$  such that (in  $GL_3(\mathfrak{B})$ )

$$E_3(\mathfrak{B}, I') \subseteq E_3(\mathfrak{A}, I).$$

Recall that  $\text{Ker}[K_1(\mathfrak{A}, I) \rightarrow K_1(\mathfrak{B})]$  is finite (see Corollary X.3.6 in [2]). Clearly  $GL_3(\mathfrak{A}, I)$  has finite index in  $GL_3(\mathfrak{B})$ , and it follows that  $E_3(\mathfrak{A}, I)$  has finite index in  $E_3(\mathfrak{B})$ . So there is a normal subgroup  $G \triangleleft E_3(\mathfrak{B})$  of finite index and contained in  $E_3(\mathfrak{A}, I)$ . Choose  $I'$  small enough so that  $e'_{ij} \in G$  for all  $r \in I'$  and  $1 \leq i, j \leq 3$  (e.g., take  $I' \subseteq e\mathfrak{B}$  where  $e = \exp(E_3(\mathfrak{B})/G)$ ). Since

$E_3(\mathfrak{B}, I')$  is the smallest normal subgroup of  $E_3(\mathfrak{B})$  containing all such  $e'_{ij}$ , we have the required relation:

$$E_3(\mathfrak{B}, I') \subseteq G \subseteq E_3(\mathfrak{A}, I).$$

*Step 2.* The exact sequence for an ideal induces an injection

$$\varprojlim_I \text{Coker}[K_2(\mathfrak{A}) \rightarrow K_2(\mathfrak{A}/I)] \rightarrow \varprojlim_I K_1(\mathfrak{A}, I)$$

(where limits are still taken over all ideals of finite index), and similarly for  $\mathfrak{B}$ . Hence, by Step 1, the map

$$\varphi_1: \varprojlim_I \text{Coker}[K_2(\mathfrak{A}) \rightarrow K_2(\mathfrak{A}/I)] \rightarrow \varprojlim_I \text{Coker}[K_2(\mathfrak{B}) \rightarrow K_2(\mathfrak{B}/I)]$$

is injective.

Recall that for any finite ring  $R$ ,  $K_2(R)$  is finite with torsion only for primes dividing  $|R|$  (see, e.g., Theorem 2.4 in [11]). In particular, for any  $I \subseteq \mathfrak{A}$  of finite index,

$$\begin{aligned} \text{Coker}[K_2(\mathfrak{A}) \rightarrow K_2(\mathfrak{A}/I)] &\cong \text{Coker}[K_2(\mathfrak{A}) \rightarrow K_2((\mathfrak{A}/I)_{(p)})] \\ &\quad \oplus \text{Coker}[K_2(\mathfrak{A}) \rightarrow K_2((\mathfrak{A}/I)[1/p])] \end{aligned}$$

(and similarly for  $\mathfrak{B}$ ). So  $\varphi_1$  splits as a corresponding sum of injections. But

$$\varprojlim_I \text{Coker}[K_2(\mathfrak{A}) \rightarrow K_2((\mathfrak{A}/I)_{(p)})] \cong \varprojlim_n \text{Coker}[K_2(\mathfrak{A}) \rightarrow K_2(\mathfrak{A}/p^n \mathfrak{A})],$$

and so the map  $\varphi$  above is injective. ■

Based on the notation in [1], we now set

$$\begin{aligned} C(\Sigma, A) &= \varprojlim_I \text{Coker}[K_2(\mathfrak{M}) \rightarrow K_2(\mathfrak{M}/I)] \\ &\cong \prod_p \varprojlim_n \text{Coker}[K_2(\mathfrak{M}) \rightarrow K_2(\mathfrak{M}/p^n \mathfrak{M})] \end{aligned}$$

for any semisimple  $\mathbb{Q}$ -algebra  $A$  and maximal  $\mathbb{Z}$ -order  $\mathfrak{M} \subseteq A$ . By Theorem 5.2 in [1],  $C(\Sigma, A)$  is finite for any  $A$ . Using this, we now get

**THEOREM 5.2.** *Let  $\pi$  be a  $p$ -group, and  $R$  the ring of integers in some finite extension  $K$  of  $\mathbb{Q}$  where  $p$  is unramified. Set*

$$C^*(\Sigma, K\pi) = C(\Sigma, K\pi)_{(p)} / \langle \{a, u\} \in C(\Sigma, K\pi) \mid a \in (\hat{R}_p)^*, u \in (\hat{R}_p \pi)^* \rangle,$$

and let  $r = rk_{\mathbb{Z}}(R)$ . Then

$$|K_2(R\pi)_{(p)}| \geq \frac{|\mathcal{U}(\pi)|^r \cdot |SK_1(R\pi)|}{|C^*(\Sigma, K\pi)|}$$

and

$$|Wh_2(\pi)_{(p)}| \geq \frac{|\mathcal{U}(\pi)| \cdot |SK_1(\mathbb{Z}\pi)|}{|H_2(\pi)| \cdot |C^*(\Sigma, \mathbb{Q}\pi)|}.$$

*Proof.* Let  $\mathfrak{M} \supseteq R\pi$  be a maximal order in  $K\pi$ ;  $Cl_1(\mathfrak{M}) = 0$  by [7]. Applying Theorem 5.1 to the pair  $R\pi \subseteq \mathfrak{M}$ , we get

$$\varprojlim_n \text{Coker}[K_2(R\pi) \rightarrow K_2(R/p^n[\pi])] = |C(\Sigma, K\pi)_{(p)}| / |Cl_1(R\pi)|. \quad (1)$$

Furthermore,

$$\varprojlim_n \text{Coker}[K_2(R\pi) \rightarrow K_2(R/p^n[\pi])] \cong \text{Coker}[K_2(R\pi) \rightarrow K_2^{\text{top}}(\hat{R}_p\pi)]$$

if  $K_2(R\pi)$  is finite, and so in any case

$$|K_2(R\pi)_{(p)}| \geq |K_2^{\text{top}}(\hat{R}_p\pi)| / |\varprojlim_n \text{Coker}[K_2(R\pi) \rightarrow K_2(R/p^n[\pi])]|. \quad (2)$$

By Proposition 0.1,  $K_2^{\text{top}}(\hat{R}_p\pi)$  maps onto  $K_2^*(\hat{R}_p\pi)$ , and symbols of the form  $\{a, u\}$  for  $a \in (\hat{R}_p)^*$  and  $u \in (\hat{R}_p\pi)^*$  clearly lie in the kernel. It follows that

$$|K_2^{\text{top}}(\hat{R}_p\pi)| \geq |K_2^*(\hat{R}_p\pi)| \cdot |\langle \{a, u\} \in C(\Sigma, K\pi)_{(p)} : a \in (\hat{R}_p)^*, u \in (\hat{R}_p\pi)^* \rangle|. \quad (3)$$

Write  $\hat{R}_p = \sum A_i$ , a sum of unramified  $p$ -rings. For each  $i$ ,

$$\mathcal{U}_0(A_i\pi) = \text{Ker}[v: \mathcal{U}(A_i\pi) \rightarrow H_2(\pi)],$$

where

$$\text{Im}(v) = \langle g \wedge h : g, h \in \pi, gh = hg \rangle = H_2^{ab}(\pi)$$

(see Section 3 in [10]). Hence by Theorem 3.6,

$$\begin{aligned} |K_2^*(A_i\pi)| &\geq |\mathcal{U}_0(A_i\pi)| \cdot |H_2(\pi)| = |A_i \otimes \mathcal{U}(\pi)| \cdot |H_2(\pi)| / |H_2^{ab}(\pi)| \\ &= |A_i \otimes \mathcal{U}(\pi)| \cdot |SK_1(A_i\pi)| \end{aligned}$$

(also applying Theorem 3 in [10]), and so

$$|K_2^*(\hat{R}_p\pi)| \geq |\hat{R}_p \otimes \mathcal{U}(\pi)| \cdot |SK_1(\hat{R}_p\pi)| = |\mathcal{U}(\pi)|^r \cdot |SK_1(\hat{R}_p\pi)|. \quad (4)$$

Combining (1), (2), (3), and (4) now gives the formula

$$|K_2(R\pi)_{(p)}| \geq |\mathcal{U}(\pi)|^r \cdot |SK_1(R\pi)| / |C^*(\Sigma, K\pi)|.$$

To obtain the second inequality, we must check that

$$\begin{aligned} & |K_2^{\text{top}}(\hat{\mathbb{Z}}_p \pi) / \langle \{-1, -1\}, \{-1, g\} : g \in \pi \rangle | \\ & \geq |K_2^*(\hat{\mathbb{Z}}_p \pi)| \cdot |\langle \{a, u\} \in C(\Sigma, \mathbb{Q}\pi)_{(p)} \rangle|. \end{aligned}$$

If  $p$  is odd, this follows directly from (3). If  $p=2$ , then the symbols  $\{-1, -1\}$  and  $\{-1, g\}$  are clearly trivial in  $K_2^*(\hat{\mathbb{Z}}_p \pi)$ , and we must check that they also map trivially to  $C(\Sigma, \mathbb{Q}\pi)$ . It suffices to check this when  $\pi$  is cyclic. But  $C(\Sigma, \mathbb{Q})=0$ , and for any  $n \geq 2$  and any primitive  $2^n$ -th root of unity  $\zeta$ ,

$$\{-1, \zeta\} = 1 \quad \text{in} \quad K_2^{\text{top}}(\hat{\mathbb{Z}}_2[\zeta]) \cong C(\Sigma, \mathbb{Q}\zeta)_{(2)}$$

(this follows easily from Theorem 15.7 in [9]). ■

For abelian  $p$ -groups, for example, more explicit lower bounds for  $|K_2(\mathbb{Z}\pi)|$  can easily be derived from those in Theorem 5.3, generalizing the formulas in [4].

## 6

We now study the groups

$$K_2^*(A/p^n[\pi]) = \varprojlim_{\tilde{\pi} \rightarrow \pi} \text{Coker}[K_2(A/p^n[\tilde{\pi}]) \rightarrow K_2(A/p^n[\pi])]$$

when  $\pi$  is a  $p$ -group,  $A$  an unramified  $p$ -ring, and  $n \geq 1$ . A description of them will be given analogous to the description of  $K_2^*(A\pi)$  in Theorem 3.6. This will then be used to show that  $K_2^*(\mathbb{F}_q\pi) = K_2(\mathbb{F}_q\pi)$  when  $\pi$  is an abelian  $p$ -group (and  $q = p^f$ ), thus providing a complete description of  $K_2(\mathbb{F}_q\pi)$  in this case.

The relationship between  $K_2^*(A/p^n[\pi])$  and  $K_2^*(A\pi)$  is described by:

**PROPOSITION 6.1.** *For any  $p$ -group  $\pi$  and unramified  $p$ -ring  $A$ , define a homomorphism*

$$\psi: \mathcal{U}(A\pi) \rightarrow K_2^*(A\pi)$$

*by setting  $\psi(h \otimes \lambda g) = \{h, \exp(p\lambda(g-1))\}$  for  $\lambda \in A$  and commuting  $g, h \in \pi$ . Then for any  $n \geq 1$ , the sequence*

$$\mathcal{U}(A\pi) \xrightarrow{p^{n-1}\psi} K_2^*(A\pi) \rightarrow K_2^*(A/p^n[\pi]) \rightarrow 0 \quad (1)$$

is exact. Furthermore, in the notation of Theorem 3.6,

$$\omega_2\psi(h \otimes \lambda g) = p\lambda(h \wedge g) = p\tilde{v}(h \otimes \lambda g) \in A \otimes H_2(\pi)$$

and (letting  $\varphi \in \text{Gal}(A/\widehat{\mathbb{Z}}_p)$  denote the Frobenius automorphism)

$$\Gamma_2^*\psi(h \otimes \lambda g) = h \otimes p\lambda g - h \otimes \varphi(\lambda) g^p \in \mathcal{U}(A\pi).$$

*Proof.* Note first that  $\psi$  is well defined: if  $g \in \pi$ ,  $\lambda \in A$ , and  $n \in \mathbb{Z}$ , then

$$\{g, \exp(p\lambda(g^n - 1))\} = 1 \quad \text{in } K_2^*(A\pi)$$

(the symbol lifts to arbitrary large  $p$ -groups  $\tilde{\pi} \rightarrow \pi$ ). Clearly,

$$\text{Im}(p^{n-1}\psi) \subseteq \text{Ker}[K_2^*(A\pi) \rightarrow K_2^*(A/p^n[\pi])].$$

To show inclusion the other way, choose an extension

$$1 \rightarrow \rho \rightarrow \tilde{\pi} \xrightarrow{\alpha} \pi \rightarrow 1$$

of  $p$ -groups such that

$$K_2^*(A\pi) = \text{Coker}[K_2(A\tilde{\pi}) \rightarrow K_2(A\pi)],$$

$$K_2^*(A/p^n[\pi]) = \text{Coker}[K_2(A/p^n[\tilde{\pi}]) \rightarrow K_2(A/p^n[\pi])].$$

Let  $\mathfrak{U} \subseteq A\tilde{\pi} \times A\tilde{\pi}$  be the pullback over  $A\pi$ , as usual, and let  $r_i: \mathfrak{U} \rightarrow A\tilde{\pi}$  denote the projections. Then  $\mathfrak{U}/p^n\mathfrak{U}$  is the pullback of  $A/p^n[\tilde{\pi}]$  with itself over  $A/p^n[\pi]$ , and so there are Mayer-Vietoris sequences

$$\begin{array}{ccccccc} 0 \rightarrow & K_2^*(A\pi) & \xrightarrow{\partial} & K_1(\mathfrak{U}) & \xrightarrow{r_1^* \oplus r_2^*} & K_1(A\tilde{\pi}) \oplus K_1(A\tilde{\pi}) \\ & \downarrow & & \downarrow & & \downarrow \\ 0 \rightarrow & K_2^*(A/p^n[\pi]) & \longrightarrow & K_1(\mathfrak{U}/p^n\mathfrak{U}) & \longrightarrow & K_1(A/p^n[\tilde{\pi}]) \oplus K_1(A/p^n[\tilde{\pi}]) \end{array}$$

Hence, for any

$$\xi \in \text{Ker}[K_2^*(A\pi) \rightarrow K_2^*(A/p^n[\pi])],$$

we have

$$\partial(\xi) \in \text{Ker}[K_1(\mathfrak{U}) \rightarrow K_1(\mathfrak{U}/p^n\mathfrak{U})] = \text{Im}[K_1(\mathfrak{U}, p^n\mathfrak{U}) \rightarrow K_1(\mathfrak{U})].$$

Write  $\partial(\xi) = [1 + p^n x]$  for some  $x \in \mathfrak{U}$ . Since  $r_2$  is split by the diagonal map  $\Delta: A\tilde{\pi} \rightarrow \mathfrak{U}$ , we may assume that  $x \in \text{Ker}(r_2)$ , and hence  $r_1(x) \in \text{Ker}(\alpha)$ .

Set

$$p^n y = \log(1 + p^n \cdot r_1(x)) \quad (y \in \text{Ker}[\alpha: A\tilde{\pi} \rightarrow A\pi]).$$

Then  $1 + p^n \cdot r_1(x) = \exp(p^n y)$ : this is clear if  $p^n > 2$ , and holds when  $p^n = 2$  since  $r_1(x) \in \text{Ker}(\alpha) \subseteq J(A\tilde{\pi})$  (and thus  $x^k \rightarrow 0$  as  $k \rightarrow \infty$ ). In other words,

$$[\exp(p^n y)] = r_{1*}(\partial(\xi)) \in K_1(A\tilde{\pi}, \text{Ker}(\alpha)). \quad (2)$$

Write  $y = \sum_{k \in \pi} y_k$ , where  $y_k \in A(\alpha^{-1}k)$  for  $k \in \pi$ . Then  $y_k \in \text{Ker}(\alpha)$  for all  $k$ , and  $\exp(p^n y_k) = \exp(p^n g y_k g^{-1})$  in  $K_1(A\tilde{\pi}, \text{Ker}(\alpha))$  for any  $g \in \tilde{\pi}$ . So after conjugating, we may assume that  $y = \sum y_{k_i}$ , where the  $k_i$  run through a set of conjugacy class representatives for  $\pi$ . Since  $[\exp(p^n y)] = 1$  in  $K_1(A\tilde{\pi})$ , we must have  $y = 0$  in  $\overline{A\tilde{\pi}}$ , and hence each  $y_{k_i}$  is a sum of terms of the form  $(hgh^{-1} - g)$  for  $g, h \in \tilde{\pi}$  such that  $[\alpha(g), \alpha(h)] = 1$  in  $\pi$ .

So we can now write

$$y = \sum_i \lambda_i (h_i g_i h_i^{-1} - g_i) \quad (\lambda_i \in A, g_i, h_i \in \tilde{\pi}, [\alpha g_i, \alpha h_i] = 1).$$

Then by (2):

$$\begin{aligned} \partial(\xi) &= \partial \left( \prod_i \{ \alpha(h_i), \exp(p^n \lambda_i (\alpha(g_i) - 1)) \} \right) \\ &= p^{n-1} \partial \psi \left( \sum_i \alpha(h_i) \otimes \lambda_i \alpha(g_i) \right). \end{aligned}$$

Since  $\partial$  is one-to-one, this shows that  $\xi \in \text{Im}(p^{n-1}\psi)$ .

Thus, sequence (1) is exact at  $K_2^*(A\pi)$ . Exactness at  $K_2^*(A/p^n[\pi])$  follows since  $K_2(A\pi)$  surjects onto  $K_2(A/p^n[\pi])$  by Lemma 1.7 in [11] (or, when  $p^n = 2$ , since the only torsion in  $K_1(A\pi, 2A\pi)$  is  $-1$  by Proposition 2 in [10]). The descriptions of  $\omega_2\psi$  and  $\Gamma_2^*\psi$  follow immediately from Propositions 3.7 and 3.8 (and the definition of  $\Gamma$  in [10]). ■

There are several ways of combining Proposition 6.1 with Theorem 3.6 to get descriptions of  $K_2^*(A/p^n[\pi])$ ; we give only one of them here. Note in particular that  $H_2(\pi)$  still injects into  $K_2^*(A/p^n[\pi])$ .

**PROPOSITION 6.2.** *For any  $p$ -group  $\pi$ , any unramified  $p$ -ring  $A$ , and any  $n \geq 1$ , there is a short exact sequence*

$$\begin{aligned} 0 \longrightarrow H_2(\pi) &\longrightarrow \frac{K_2^*(A/p^n[\pi])}{\text{Im}(\bar{\eta})} \\ &\xrightarrow{\Gamma_2^*} \frac{\mathcal{U}_0(A\pi)}{\langle h \otimes p^n \lambda g - h \otimes p^{n-1} \varphi(\lambda) g^p \rangle} \longrightarrow 0. \end{aligned}$$

Here  $\bar{\Gamma}_2^*$  is the reduction of  $\Gamma_2^*$ , and  $\bar{\eta}$  denotes the composite

$$\bar{\eta}: H_3(\pi) \xrightarrow{\eta_{A\pi}} K_2^*(A\pi) \rightarrow K_2^*(A/p^n[\pi]).$$

*Proof.* Let  $\psi: \mathcal{U}(A\pi) \rightarrow K_2^*(A\pi)$  be the map defined above. By Proposition 6.1 and Theorem 3.6,  $(\omega_2, \Gamma_2^*)$  induces an injection

$$\frac{K_2^*(A/p^n[\pi])}{\text{Im}(\bar{\eta})} \xrightarrow{\quad} \frac{(A \otimes H_2(\pi)) \oplus \mathcal{U}(A\pi)}{p^{n-1}(\omega_2\psi, \Gamma_2^*\psi)(\mathcal{U}(A\pi))}.$$

Then (using Theorem 3.6(1) again)

$$\begin{aligned} \text{Ker} \left[ \bar{\Gamma}_2^*: \frac{K_2(A/p^n[\pi])}{\text{Im}(\bar{\eta})} \rightarrow \frac{\mathcal{U}(A\pi)}{p^{n-1}\Gamma_2^*\psi(\mathcal{U}(A\pi))} \right] \\ \cong \frac{\omega_2(\text{Ker}(\Gamma_2^*))}{\omega_2(\text{Ker}(\Gamma_2^*) \cap \text{Im}(\psi))} \\ = \frac{\text{Ker}[(1-\varphi) \otimes 1: A \otimes H_2(\pi) \rightarrow A \otimes H_2(\pi)]}{\omega_2\psi(\text{Ker}(\Gamma_2^*\psi))} = \frac{H_2(\pi)}{\omega_2\psi(\text{Ker}(\Gamma_2^*\psi))}. \end{aligned}$$

Furthermore (see Theorem 3.6(2))

$$\text{Im}(\bar{\Gamma}_2^*) = \frac{\text{Im}(\Gamma_2^*)}{p^{n-1}\Gamma_2^*\psi(\mathcal{U}(A\pi))} = \frac{\mathcal{U}_0(A\pi)}{\langle h \otimes p^n \lambda g - h \otimes p^{n-1} \varphi(\lambda) g^p \rangle}.$$

So we will be done upon showing that  $\text{Ker}(\Gamma_2^*\psi) \subseteq \text{Ker}(\omega_2\psi)$ .

Define

$$\Phi: \mathcal{U}(A\pi) \rightarrow \mathcal{U}(A\pi)$$

by setting  $\Phi(h \otimes \lambda g) = h \otimes \varphi(\lambda) g^p$ . Then

$$\Gamma_2^*\psi(x) = px - \Phi(x)$$

for any  $x \in \mathcal{U}(A\pi)$ . Fix

$$x = \sum_{i=1}^k x_i \in \text{Ker}(\Gamma_2^*\psi),$$

where for each  $i$ ,  $x_i$  is the sum of terms  $h \otimes \lambda g$  with  $|g| = p^i$  (note that  $h \otimes \lambda 1 = 0$  in  $\mathcal{U}(A\pi)$ ). Then

$$\begin{aligned} 0 = \Gamma_2^*\psi(x) &= \sum_{i=1}^k (px_i - \Phi(x_i)) \\ &= px_k + \sum_{i=1}^{k-1} (px_i - \Phi(x_{i+1})), \end{aligned}$$

so that  $px_k = 0$  and  $px_i = \Phi(x_{i+1})$  for all  $i < k$ . Hence, for all  $i$ ,

$$\tilde{v}(px_i) = \tilde{v}(\Phi(x_{i+1})) = (\varphi \otimes 1)(\tilde{v}(px_{i+1})) \in A \otimes H_2(\pi),$$

and by induction

$$\tilde{v}(px_i) = (\varphi^{k-i} \otimes 1)(\tilde{v}(px_k)) = 0.$$

Thus,  $\omega_2 \psi(x) = p\tilde{v}(x) = 0$ , and we are done. ■

Note that the short exact sequence of Proposition 6.2 is *not* split by  $\omega_2$  when  $A = \hat{\mathbb{Z}}_p$ , since  $\omega_2 \psi \neq 0$  in general.

For the rest of this section, we restrict attention to abelian  $p$ -groups. We will show that  $K_2(\mathbb{F}_q \pi) = K_2^*(\mathbb{F}_q \pi)$  for abelian  $\pi$  (and  $q = p^f$ ) by first computing  $|K_2^*(\mathbb{F}_q \pi)|$ ; and then showing that  $|K_2(\mathbb{F}_q \pi)|$  can be at most as large.

For any abelian  $p$ -group  $\pi$  and any  $i \geq 1$ , we write

$$p^i\text{-rk}(\pi) = \dim_{\mathbb{F}_p}(\pi^{p^{i-1}}/\pi^{p^i}),$$

i.e., the number of cyclic summands in the decomposition of  $\pi$  of order at least  $p^i$ . Also,  $\text{ord}_p(|\pi|)$  is used to denote the number  $n$  such that  $|\pi| = p^n$ .

**PROPOSITION 6.3.** *For any abelian  $p$ -group  $\pi$ , any unramified  $p$ -ring  $A$  with  $f = [A : \hat{\mathbb{Z}}_p]$ , and any  $n \geq 1$ ,*

$$|K_2^*(A/p^n[\pi])| = |\mathcal{U}(\pi) \otimes \mathbb{Z}/p^n|^f.$$

If  $\exp(\pi) = p^e$  and  $r_i = p^i\text{-rk}(\pi)$  ( $1 \leq i \leq e$ ), then

$$\begin{aligned} \text{ord}_p |K_2^*(A/p[\pi])| \\ = f[(r_1 - 1)|\pi| - (r_1 - r_2)|\pi^p| - \cdots - (r_{e-1} - r_e)|\pi^{p^{e-1}}| - (r_e - 1)]. \end{aligned}$$

*Proof.* *Step 1.* We first claim that

$$S_\pi(g) = \{h \in \pi : h^{p^m} \in \langle g \rangle\} \quad (1)$$

for any  $m \geq 0$  and  $g \in \pi^{p^m} - \pi^{p^{m+1}}$ . Recall that  $S_\pi(g)$  is the subgroup generated by all  $h \in \pi$  such that  $g \in \langle h \rangle$ , then  $\langle h^{p^m}, g \rangle$  is cyclic for such  $h$ , and  $h^{p^m} \in \langle g \rangle$  since  $g \notin \langle h^{p^{m+1}} \rangle$  by assumption. Conversely, if  $h^{p^m} = g^a$  and  $g = g_0^{p^m}$  for some  $a \in \mathbb{Z}$  and  $g_0 \in \pi$ , then  $(hg_0^{1-a})^{p^m} = g$ , and so

$$h = g^{a-1}(hg_0^{1-a}) \in S_\pi(g).$$

*Step 2.* Again, consider the map

$$\Phi: \mathcal{U}(A\pi) \rightarrow \mathcal{U}(A\pi) \quad (\Phi(h \otimes \lambda g) = h \otimes \varphi(\lambda) g^p).$$



Since  $\pi$  is abelian,

$$\mathcal{U}(A\pi) \cong \sum_{g \in \pi} (\pi/S_\pi(g)) \otimes A(g).$$

So  ${}_p\mathcal{U}(A\pi)$  is generated by elements of the form  $h \otimes \lambda g$ , where  $h^p \in S_\pi(g)$ . Fix such an element, and assume  $g \in \pi^{p^m} - \pi^{p^{m+1}}$  ( $m \geq 0$ ). Then  $h^{p^{m+1}} = g^a$  for some  $a$ , by (1), and  $p \mid a$  since  $g \notin \pi^{p^{m+1}}$ . So  $h^{p^{m+1}} \in \langle g^p \rangle$ ,  $g^p \in \pi^{p^n} - \pi^{p^{n+1}}$  for some  $n \geq m+1$ , and  $h \in S_\pi(g^p)$  by (1) again. In other words,  $\Phi(h \otimes \lambda g) = 0$ , and we have shown that

$${}_p\mathcal{U}(A\pi) \subseteq \text{Ker}(\Phi). \quad (2)$$

We now claim that

$$\text{Ker}(\Gamma_2^*\psi) = {}_p\mathcal{U}(A\pi). \quad (3)$$

Since  $\Gamma_2^*\psi(x) = px - \Phi(x)$  by Proposition 6.1,  $\text{Ker}(\Gamma_2^*\psi)$  contains all  $p$ -torsion by (2). Conversely, consider any

$$x = \sum_{i=1}^k x_i \in \text{Ker}(\Gamma_2^*\psi),$$

where each  $x_i$  is a sum of terms  $h \otimes \lambda g$  with  $|g| = p^i$ . Then

$$\Gamma_2^*\psi(x) = px_k + \sum_{i=1}^{k-1} (px_i - \Phi(x_{i+1})) = 0,$$

so  $px_k = 0$  and  $px_i = \Phi(x_{i+1})$  for all  $i < k$ . By (2) and induction downwards from  $k$ , we get  $px_i = 0$  for all  $i$ , and hence  $x \in {}_p\mathcal{U}(A\pi)$ .

Then, for all  $n \geq 1$ ,

$$\text{Ker}(p^{n-1}\Gamma_2^*\psi) = {}_{p^n}\mathcal{U}(A\pi).$$

Since  $\eta_{A\pi} = 0$  by Theorem 4.2, Proposition 6.2 implies

$$\begin{aligned} |K_2^*(A/p^n[\pi])| &= |H_2(\pi)| \cdot |\mathcal{U}_0(A\pi)/\text{Im}(p^{n-1}\Gamma_2^*\psi)| \\ &= |\mathcal{U}(A\pi)/\text{Im}(p^{n-1}\Gamma_2^*\psi)| = |\text{Ker}(p^{n-1}\Gamma_2^*\psi)| \quad (4) \\ &= |{}_p\mathcal{U}(A\pi)| = |\mathcal{U}(A\pi) \otimes \mathbb{Z}/p^n| = |\mathcal{U}(\pi) \otimes \mathbb{Z}/p^n|^f \end{aligned}$$

(recall that  $\mathcal{U}(A\pi) \cong A \otimes \mathcal{U}(\pi)$ , and  $f = [A: \widehat{\mathbb{Z}}_p]$ ).

Step 3. It remains only to compute the order of

$$\mathcal{U}(\pi) \otimes \mathbb{Z}/p \cong \sum_{g \in \pi} (\pi/S_\pi(g)) \otimes \mathbb{Z}/p.$$

If  $g \in \pi^{p^m} - \pi^{p^{m+1}}$  for any  $0 \leq m < e$  ( $p^e = \exp(\pi)$ ), and  $g = g_0^{p^m}$ , then

$$S_\pi(g) = \{h \in \pi: h^{p^m} \in \langle g \rangle\} = \langle g_0; x: x^{p^m} = 1 \rangle$$

by (1). In other words,  $\pi/S_\pi(g)$  has rank one less than the number of cyclic summands of order at least  $\mathbb{Z}/p^{m+1}$ :

$$\text{ord}_p |(\pi/S_\pi(g)) \otimes \mathbb{Z}/p| = r_{m+1} - 1,$$

where  $r_{m+1} = p^{m+1} - rk(\pi)$ . So

$$\begin{aligned} \text{ord}_p |\mathcal{U}(\pi) \otimes \mathbb{Z}/p| &= \sum_{m=0}^{e-1} (r_{m+1} - 1)(|\pi^{p^m}| - |\pi^{p^{m+1}}|) \\ &= r_1 \cdot |\pi| - \sum_{m=1}^{e-1} (r_m - r_{m+1}) |\pi^{p^m}| - r_e |\pi^{p^e}| - (|\pi| - 1) \\ &= (r_1 - 1) |\pi| - \sum_{m=1}^{e-1} (r_m - r_{m+1}) |\pi^{p^m}| - (r_e - 1). \end{aligned}$$

Combining this with (4) gives the formula for  $|K_2^*(A/p[\pi])|$ . ■

The formula

$$|\mathcal{U}(A\pi)/\text{Im}(\Gamma_2^* \psi)| = |\mathcal{U}(A\pi) \otimes \mathbb{Z}/p|$$

does, in fact, fail in general for non-abelian  $p$ -groups.

The techniques in [19] and [4] will now be used to find upper bounds for  $|K_2(A/p[\pi])|$ . The following symbol identities are in essence contained in Sections II.3 and III.4 of [19].

**PROPOSITION 6.4.** *Let  $p$  be a prime, and let  $R$  be a commutative ring of characteristic  $p$ . Fix  $a \in R^*$ .*

(i) *If  $t \in R$  is such that  $t^{mp^k} = 0$  for some  $k \geq 1$  and  $p \nmid m$ , then*

$$\{1 + t, 1 + a^{p^k} t^{mp^k - 1}\} = 1 \quad \text{in } K_2(R).$$

(ii) *If  $t \in R$  is such that  $t^{mp^k + 1} = 0$  for some  $k \geq 1$  and  $p \nmid m$ , and  $l < k$ , then*

$$\{a, 1 + a^{p^l} t^{mp^k}\} = 1 \quad \text{in } K_2(R).$$

*Proof.* Set  $A = \mathbb{F}_p[a, a^{-1}]$  and  $A_\infty = A[[t]]$ . Then  $A[t]$  and  $A_\infty$  are regular rings by the Hilbert syzygy theorem [2, Sect. XII.2], and so by

localization and devissage results in [23, Theorem 4 and 5], there are exact sequences

$$\begin{array}{ccccc} K_2(A) & \longrightarrow & K_2(A[t]) & \xrightarrow{i_1} & K_2(A[t, t^{-1}]) \\ \downarrow = & & \downarrow & & \downarrow \\ K_2(A) & \longrightarrow & K_2(A_\infty) & \xrightarrow{i_2} & K_2(A_\infty[t^{-1}]) \end{array}$$

Since  $i_1$  is injective by the corollary to Theorem 8 in [23],  $i_2$  must also be injective.

Fix integers  $k, m \geq 1$  and  $0 \leq l \leq k$ . Then relations of the form  $\{u, 1-u\} = 1$  in  $K_2(A_\infty[t^{-1}])$  give

$$\begin{aligned} & \{(-1)^{mp^{k-l}+1}a, 1+a^{p^l}t^{mp^k}\} \\ &= \{(-1)^{mp^{k-l}+1}a, 1+at^{mp^{k-l}}\}^{p^l} \\ &= \{(-t)^{mp^{k-l}}, 1+at^{mp^{k-l}}\}^{-p^l} = \{(-t)^{mp^k}, 1+at^{mp^{k-l}}\}^{-1} \\ &= \{-t, 1+a^{p^k}t^{mp^{2k-l}}\}^{-m} = \{-t, (1+t)(1+a^{p^k}t^{mp^{2k-l}})\}^{-m} \\ &= \{1+(1+t)a^{p^k}t^{mp^{2k-l}-1}, (1+t)(1+a^{p^k}t^{mp^{2k-l}})\}^m. \end{aligned}$$

So by injectivity of  $i_2$ ,

$$\begin{aligned} & \{1+(1+t)a^{p^k}t^{mp^{2k-l}-1}, (1+t)(1+a^{p^k}t^{mp^{2k-l}})\}^m \\ &= \{(-1)^{m+1}a, 1+a^{p^l}t^{mp^k}\} \quad \text{in } K_2(A_\infty). \end{aligned} \quad (1)$$

Now consider the individual formulas to be shown.

(i) It suffices to prove this when  $R = A_\infty/(t^{mp^k})$ . When  $l=k$ , (1) reduces to the formula

$$\{1+a^{p^k}t^{mp^{k-1}}, 1+t\}^m = 1 \quad \text{in } K_2(R).$$

Since  $p \nmid m$  and  $(1+a^{p^k}t^{mp^{k-1}})^p = 1$  in  $R$ , this gives

$$\{1+t, 1+a^{p^k}t^{mp^{k-1}}\} = 1.$$

(ii) It suffices to show this when  $R = A_\infty/(t^{mp^k+1})$ . Since  $k > l \geq 0$ ,  $t^{mp^{2k-l}-1} = 0$  in  $R$ . So by (1),

$$\{(-1)^{m+1}a, 1+a^{p^l}t^{mp^k}\} = 1 \quad \text{in } K_2(R).$$

But  $\{-1, 1+a^{p^l}t^{mp^k}\} = 1$  since  $(1+a^{p^l}t^{mp^k})^p = 1$  in  $R$  (and  $-1 = 1$  if  $p=2$ ). ■

Now consider the groups

$$\Phi_n(R) = \text{Ker}[K_2(R[t]/(t^{n+1})) \rightarrow K_2(R[t]/(t^n))],$$

defined for any ring  $R$  and  $n \geq 1$ .

LEMMA 6.5. *Let  $\pi$  be an abelian  $p$ -group, and set  $r = rk(\pi)$ . Then, for any  $q = p^f$  and any  $n \geq 1$ ,*

$$\begin{aligned} \text{ord}_p |\Phi_n(\mathbb{F}_q \pi)| &\leq fr |\pi| && \text{if } n \not\equiv -1, 0 \pmod{p}, \\ &\leq fr |\pi| + f(|\pi| - |\pi^{p^k}|) && \text{if } n = mp^k - 1, k \geq 1, p \nmid m, \\ &\leq fr |\pi| - f(|\pi| - |\pi^{p^k}|) && \text{if } n = mp^k, k \geq 1, p \nmid m. \end{aligned}$$

*Proof.* Set  $R = \mathbb{F}_q \pi[t]/(t^{n+1})$ . Let  $\{g_1, \dots, g_r\}$  be generators for  $\pi$ , and let

$$J = \langle g_1 - 1, g_2 - 1, \dots, g_r - 1, t \rangle$$

denote the Jacobson radical of  $R$ . For convenience, set

$$K_2(R, I) = \text{Ker}[K_2(R) \rightarrow K_2(R/I)]$$

for any ideal  $I \subseteq R$ . By Proposition 1.1 in [17],

$$\begin{aligned} \Phi_n(\mathbb{F}_q \pi) &= K_2(R, t^n) \\ &= \langle \{g_i, 1 + yt^n\}, \{1 + t, 1 + yt^n\} \mid 1 \leq i \leq r, y \in \mathbb{F}_q \pi \rangle. \end{aligned} \quad (1)$$

We now consider the individual cases.

(i) Assume  $n \not\equiv -1, 0 \pmod{p}$ . For all  $y \in \mathbb{F}_q \pi$ ,

$$\{1 + t, 1 + yt^n\}^{n+1} = 1 \quad \text{in } K_2(R),$$

by [20, Corollary 1.5b]. So  $\{1 + t, 1 + yt^n\} = 1$  since  $(1 + yt^n)^p = 1$  in  $R$ , and  $\Phi_n(\mathbb{F}_q \pi)$  is generated by all  $\{g_i, 1 + yt^n\}$ . Hence,

$$\text{ord}_p |\Phi_n(\mathbb{F}_q \pi)| \leq r \cdot rk_{\mathbb{F}_p}(\mathbb{F}_q \pi) = fr |\pi|.$$

(ii) If  $n = mp^k - 1$  where  $k \geq 1$  and  $p \nmid m$ , then by Proposition 6.4,

$$\{1 + t, 1 + \lambda a^{p^k} t^n\} = 1$$

for any  $\lambda \in \mathbb{F}_q$  and  $a \in \pi$ . So by (1),  $\Phi_n(\mathbb{F}_q \pi)$  is generated by all

$$\{g_i, 1 + yt^n\} \quad (1 \leq i \leq r, y \in \mathbb{F}_q \pi)$$

and

$$\{1+t, 1+\lambda a t^n\} \quad (\lambda \in \mathbb{F}_q, a \in \pi - \pi^{p^k}).$$

It follows that

$$\text{ord}_p |\Phi_n(\mathbb{F}_q \pi)| \leq f r |\pi| + f(|\pi| - |\pi^{p^k}|).$$

(iii) Finally, assume  $n = m p^k$  where  $k \geq 1$  and  $p \nmid m$ . Then

$$\{1+t, 1+y t^n\} = 1 \quad \text{in } K_2(R),$$

for all  $y \in \mathbb{F}_q \pi$ , by the same argument as in (i). So

$$\Phi_n(\mathbb{F}_q \pi) = \langle \{g_i, 1 + \lambda a t^n\} : 1 \leq i \leq r, \lambda \in \mathbb{F}_q, a \in \pi \rangle. \quad (2)$$

By Proposition 6.4(ii),

$$\{a, 1 + \lambda a^{p^l} t^n\} = 1 \quad \text{in } K_2(R),$$

for any  $a \in \pi$ ,  $\lambda \in \mathbb{F}_q$ , and  $l < k$ . This provides  $f$  relations among the elements of (2) for each  $a \in \pi - \pi^{p^k}$ , and so

$$\text{ord}_p |\Phi_n(\mathbb{F}_q \pi)| \leq f r |\pi| - f(|\pi| - |\pi^{p^k}|). \quad \blacksquare$$

Finally, Lemma 6.5 can be applied to show:

**THEOREM 6.6.** *For any abelian  $p$ -group  $\pi$  and any  $q = p^f$ ,*

$$K_2(\mathbb{F}_q \pi) = K_2^*(\mathbb{F}_q \pi).$$

*In particular, if  $A$  is the unramified  $p$ -ring of degree  $f$  over  $\widehat{\mathbb{Z}}_p$ , there is a short exact sequence*

$$0 \rightarrow H_2(\pi) \rightarrow K_2(\mathbb{F}_q \pi) \rightarrow \mathcal{U}_0(A\pi) / \langle h \otimes p \lambda g - h \otimes \varphi(\lambda) g^p \rangle \rightarrow 0, \quad (1)$$

*which is (at least non-canonically) split.*

*Proof.* Fix  $\pi$ , and write  $\pi = \bar{\pi} \times \mathbb{Z}/p^n$ , where  $\bar{\pi}$  is a product of cyclic groups of order at least  $p^n$ . Let  $g \in \pi$  generate  $\mathbb{Z}/p^n$ , and set  $t = g - 1$ . Then

$$\mathbb{F}_q \pi \cong \mathbb{F}_q \bar{\pi} [g] / (g^{p^n} - 1) \cong \mathbb{F}_q \bar{\pi} [t] / (t^{p^n}).$$

Applying Lemma 6.5, we get (where  $r = rk(\bar{\pi})$ )

$$\begin{aligned} \text{ord}_p |K_2(\mathbb{F}_q \pi)| &= \text{ord}_p |K_2(\mathbb{F}_q \bar{\pi})| + \sum_{i=1}^{p^n-1} \text{ord}_p |\Phi_i(\mathbb{F}_q \bar{\pi})| \\ &\leq \text{ord}_p |K_2(\mathbb{F}_q \bar{\pi})| + (p^n - 1) f r |\bar{\pi}| + f(|\bar{\pi}| - |\bar{\pi}^{p^n}|). \end{aligned}$$

In particular,  $K_2(\mathbb{F}_q \pi) = 1$  if  $\bar{\pi} = 1$ .

If  $\bar{\pi} \neq 1$ , set

$$p^e = \exp(\bar{\pi}) = \exp(\pi), \quad r_i = p^i \cdot \text{rk}(\bar{\pi}).$$

Then  $r_1 = r_2 = \dots = r_n = r$ , and  $p^i \cdot \text{rk}(\pi) = r + 1$  for  $i \leq n$ . We may assume inductively that  $K_2(\mathbb{F}_q \bar{\pi}) = K_2^*(\mathbb{F}_q \bar{\pi})$ . So by Proposition 6.3,

$$\begin{aligned} \text{ord}_p |K_2(\mathbb{F}_q \pi)| &\leq \text{ord}_p |K_2(\mathbb{F}_q \bar{\pi})| + (p^n - 1) f r |\bar{\pi}| + f(|\bar{\pi}| - |\bar{\pi}^{p^n}|) \\ &= f[(r-1)|\bar{\pi}| - (r_n - r_{n+1})|\bar{\pi}^{p^n}| - \dots - (r_{e-1} - r_e)|\bar{\pi}^{p^{e-1}}| - (r_e - 1)] \\ &\quad + f r(|\pi| - |\bar{\pi}|) + f|\bar{\pi}| - f|\bar{\pi}^{p^n}| \\ &= f[r|\pi| - (r_n + 1 - r_{n+1})|\bar{\pi}^{p^n}| - \dots - (r_{e-1} - r_e)|\bar{\pi}^{p^{e-1}}| - (r_e - 1)] \\ &= f[r|\pi| - r_e] \quad \text{if } e > n, \\ &= f[r|\pi| - r_e] \quad \text{if } e = n. \end{aligned}$$

In both cases, we get  $|K_2(\mathbb{F}_q \pi)| \leq |K_2^*(\mathbb{F}_q \pi)|$  (applying Proposition 6.3 again), and the two groups are thus equal. The short exact sequence (1) now follows from Proposition 6.2 (and Theorem 4.2).

The construction of a splitting map  $K_2(\mathbb{F}_q \pi) \rightarrow H_2(\pi)$  for (1) is easily reduced to the case where  $\pi \cong \mathbb{Z}/p^n \times \mathbb{Z}/p^n$  ( $n \geq 1$ ). By Proposition 6.1 and Theorems 3.6 and 4.2,  $K_2^*(\mathbb{F}_q \pi)$  is a subquotient of  $(A \otimes H_2(\pi)) \oplus (\pi \otimes A\pi)$ , and hence in this case has exponent  $p^n$ . But any inclusion of  $H_2(\pi) \cong \mathbb{Z}/p^n$  into a finite abelian group of exponent  $p^n$  splits. ■

Theorem 6.6 generalizes automatically to give a description of  $K_2(F\pi)$  for any finite field  $F$  and any finite abelian group  $\pi$ :

**THEOREM 6.7.** *Let  $F$  be any finite field, and let  $\pi$  be any finite abelian group. Set  $p = \text{char}(F)$ , and let  $A$  be the unramified  $p$ -ring such that  $F \cong A/p$  (i.e.,  $[A: \hat{\mathbb{Z}}_p] = [F: \mathbb{F}_p]$ ). Let  $\varphi$  be the Frobenius automorphism of  $A$ . Then there is a (non-canonical) isomorphism*

$$K_2(F\pi) \cong \frac{\pi \otimes A\pi}{\langle g \otimes \lambda g: g \in \pi, \lambda \in A \rangle \cdot \langle h \otimes (p\lambda g - \varphi(\lambda) g^p): g, h \in \pi, \lambda \in A \rangle}.$$

*Proof.* Assume first that  $\pi$  is a  $p$ -group. Consider the short exact sequence

$$\begin{aligned} 0 &\rightarrow \frac{\mathcal{U}_0(A\pi)}{\langle h \otimes (p\lambda g - \varphi(\lambda) g^p) \rangle} \\ &\rightarrow \frac{\mathcal{U}(A\pi)}{\langle h \otimes (p\lambda g - \varphi(\lambda) g^p) \rangle} \xrightarrow{\nu} H_2(\pi) \rightarrow 0. \end{aligned}$$

Write  $\pi = \langle g_1 \rangle \times \cdots \times \langle g_k \rangle$ , where  $|g_1| \leq \cdots \leq |g_k|$ . Fix  $\lambda_0 \in A$  such that  $\text{Tr}(\lambda_0) = 1$ . Then  $v$  is split by the map which sends  $g_i \wedge g_j$  to  $g_i \otimes \lambda_0 g_j$  for each  $1 \leq i < j \leq k$ . Together with Theorem 6.6, this shows that

$$\begin{aligned} K_2(F\pi) &\cong \frac{\mathcal{U}(A\pi)}{\langle h \otimes (p\lambda g - \varphi(\lambda) g^p) \rangle} \\ &\cong \frac{\pi \otimes A\pi}{\langle g \otimes \lambda g^n \rangle \cdot \langle h \otimes (p\lambda g - \varphi(\lambda) g^p) \rangle} \\ &= \frac{\pi \otimes A\pi}{\langle g \otimes \lambda g \rangle \cdot \langle h \otimes (p\lambda g - \varphi(\lambda) g^p) \rangle} \end{aligned} \quad (1)$$

in the  $p$ -group case. (To see the last step, note that

$$g \otimes \lambda g^{p^k} \equiv g \otimes p^k \varphi^{-k}(\lambda) g \pmod{\langle h \otimes (p\lambda g - \varphi(\lambda) g^p) \rangle}$$

for any  $k \geq 1$ .)

Now let  $\pi$  be an arbitrary finite abelian group, and write  $\pi = \sigma \times \bar{\pi}$ , where  $\bar{\pi}$  is a  $p$ -group and  $p \nmid |\sigma|$ . Let  $K \supseteq A$  be the quotient field. Then  $K\sigma \cong \sum_{i=1}^s K_i$  for some fields  $K_i$  which are unramified over  $\widehat{\mathbb{Q}}_p$ . Let  $A_i \subseteq K_i$  be the rings of integers and set  $F_i = A_i/p$ . Let

$$f: A\sigma \xrightarrow{\cong} \sum_{i=1}^s A_i, \quad f': A\pi \xrightarrow{\cong} \sum_{i=1}^s A_i \bar{\pi},$$

and

$$f'': \pi \otimes A\pi \cong \bar{\pi} \otimes A\pi \xrightarrow{\cong} \sum_{i=1}^s \bar{\pi} \otimes A_i \bar{\pi}$$

be the induced isomorphisms. Also,  $K_2(F\pi) \cong \sum_{i=1}^s K_2(F_i \bar{\pi})$ , and so by (1) we will be done upon showing that

$$f''(\langle g \otimes \lambda g: g \in \pi, \lambda \in A \rangle) = \sum_{i=1}^s \langle g \otimes \lambda g: g \in \bar{\pi}, \lambda \in A_i \rangle \quad (2)$$

and  $(\varphi_i \in \text{Gal}(K_i/\widehat{\mathbb{Q}}_p)$  denotes the Frobenius automorphism)

$$f'(\langle p\lambda g - \varphi(\lambda) g^p \rangle) = \sum_{i=1}^s \langle p\lambda g - \varphi_i(\lambda) g^p \rangle. \quad (3)$$

That (2) holds is clear. That (3) holds will follow once we check that for any  $\lambda \in A$  and  $g \in \sigma$ ,

$$(\Sigma \varphi_i) \circ f(\lambda g) = f(\varphi(\lambda) g^p). \quad (4)$$

But  $f^{-1}(\Sigma\varphi_i)f$  and  $(\lambda g \mapsto \varphi(\lambda)g^p)$  are both automorphisms of  $A\sigma$ , and both congruent mod  $p$  to the map  $x \rightarrow x^p$ . Two ring automorphisms of  $A\sigma \cong A_i$  which are congruent mod  $p$  must agree on roots of unity of order prime to  $p$ , and hence must be equal. So (4) holds, and we are done. ■

In fact, Theorem 6.6, together with the results and techniques in [11] and [22], can be used to give a description of  $K_2(F\pi)$  when  $F$  is a finite field of characteristic  $p$  and  $\pi$  is any finite group with abelian  $p$ -Sylow subgroup.

Theorem 6.6 also give rise to the following obvious conjecture:

*Conjecture 6.8.* For any  $p$ -group  $\pi$  and any  $q = p^f$ ,

$$K_2(\mathbb{F}_q\pi) = K_2^*(\mathbb{F}_q\pi).$$

## 7

Our original motivation for studying  $K_2(\hat{\mathbb{Z}}_p\pi)$  was its appearance in localization sequences for computing  $Cl_1(\mathbb{Z}\pi)$  and  $SK_1(\mathbb{Z}\pi)$ . Theorem 3.6, while not enough to give a complete algorithm for computing  $Cl_1(\mathbb{Z}\pi)$ , does motivate a conjecture for such an algorithm (for computing odd torsion in  $Cl_1(\mathbb{Z}\pi)$ , at least), as well as giving upper and lower bounds for  $Cl_1(\mathbb{Z}\pi)$ .

Fix an odd prime  $p$  and a  $p$ -group  $\pi$ . For each (complex) irreducible character  $\chi$  of  $\pi$ , let  $\hat{V}_\chi$  be the corresponding irreducible  $\mathbb{C}\pi$ -representation. By [24, Sects. 2 and 3],  $\mathbb{Q}\chi$  has Schur index  $m_\chi = 1$ , and  $\mathbb{Q}\chi$  (the extension of  $\mathbb{Q}$  by all  $\chi(g)$ ) is isomorphic to the cyclotomic field  $\mathbb{Q}\zeta_{p^n}$  for some  $n$ . Hence,  $\hat{V}_\chi \cong \mathbb{C} \otimes_{\mathbb{Q}\chi} V_\chi$  for some irreducible  $\mathbb{Q}\chi[\pi]$ -module  $V_\chi$ , and the simple summand  $A_\chi$  of  $\mathbb{Q}\pi$  corresponding to  $\chi$  has the form

$$A_\chi \cong \text{End}_{\mathbb{Q}\chi}(V_\chi) \cong M_{r_\chi}(\mathbb{Q}\chi) \quad (r_\chi = \dim_{\mathbb{Q}\chi}(V_\chi)).$$

For shortness of notation, we let  $\mathbb{Z}\chi$  and  $\hat{\mathbb{Z}}_p[\chi]$  denote the rings of integers in  $\mathbb{Q}\chi$  and  $\hat{\mathbb{Q}}_p[\chi]$ , respectively.

For each  $\chi$ , let  $(\mu_{\mathbb{Q}\chi})_p$  denote the group of  $p$ -power roots of unity in  $\mathbb{Q}\chi$ . For any set  $T$  of irreducible characters of  $\pi$ , homomorphisms

$$\varphi_T = \prod_{\chi \in T} \varphi_\chi: K_2^{\text{top}}(\hat{\mathbb{Z}}_p\pi) \rightarrow \prod_{\chi \in T} K_2^{\text{top}}(\hat{\mathbb{Z}}_p[\chi]),$$

$$\psi_T = \prod_{\chi \in T} \psi_\chi: H_1(\pi; \hat{\mathbb{Z}}_p\pi) \rightarrow \prod_{\chi \in T} (\mu_{\mathbb{Q}\chi})_p,$$

and

$$\lambda_T = \prod_{\chi \in T} \lambda_\chi: \prod_{\chi \in T} K_2^{\text{top}}(\hat{\mathbb{Z}}_p[\chi]) \xrightarrow{\sim} \prod_{\chi \in T} (\mu_{\mathbb{Q}\chi})_p$$



are defined as follows. Fix  $\chi \in T$ , and let  $M_\chi \subseteq V_\chi$  be a  $\pi$ -invariant  $\mathbb{Z}\chi$ -lattice. Let  $\varphi_\chi$  be the composite

$$\begin{aligned}\varphi_\chi: K_2^{\text{top}}(\widehat{\mathbb{Z}}_p\pi) &\rightarrow K_2^{\text{top}}(\widehat{\mathbb{Z}}_p \otimes \text{End}_{\mathbb{Z}\chi}(M_\chi)) \\ &\cong K_2^{\text{top}}(M_{r_\chi}(\widehat{\mathbb{Z}}_p[\chi])) \cong K^{\text{top}}(\widehat{\mathbb{Z}}_p[\chi]).\end{aligned}$$

For any commuting elements  $g, h \in \pi$ , set

$$\psi_\chi(g \otimes h) = \det_{\mathbb{Q}\chi}(g, V_\chi^h) \in (\mu_{\mathbb{Q}\chi})_p,$$

where  $V_\chi^h \subseteq V_\chi$  is the subspace of elements fixed by  $h$ . Finally,  $\lambda_\chi$  denotes the norm residue symbol map  $\lambda_\chi(\{a, b\}) = (a, b)_v$  (and is an isomorphism by [20, Theorem 5.1]).

**PROPOSITION 7.1.** *For any irreducible character  $\chi$  of  $\pi$ , and any commuting  $g, h \in \pi$ ,*

$$\psi_\chi(g \otimes h) = \lambda_\chi \circ \varphi_\chi(\{g, \Gamma^{-1}(\tfrac{1}{2}(h + h^{-1}))\}). \quad (1)$$

For any  $g \in \pi$  and  $u \in (\widehat{\mathbb{Z}}_p[Z_\pi(g)])^*$ ,

$$\psi_\chi(g \otimes \Gamma(u)) = \lambda_\chi \circ \varphi_\chi(\{g, u\}). \quad (2)$$

*Proof.* Set  $q = p^k = \exp(\pi)$ . Let

$$A_\chi \cong \text{End}_{\mathbb{Q}\chi}(V_\chi) \cong M_r(\mathbb{Q}\chi) \subseteq M_r(\mathbb{Q}\zeta_q) \quad (r = r_\chi = \dim_{\mathbb{Q}\chi}(V_\chi))$$

be the simple summand of  $\mathbb{Q}\pi$  corresponding to  $\chi$ . If  $g, h \in \pi$  and  $[g, h] = 1$ , then let  $\alpha(g)$  and  $\alpha(h)$  denote their images in  $M_r(\mathbb{Q}\chi)$ . Since  $\langle g, h \rangle$  is an abelian group of exponent dividing  $q$ , the matrices  $\alpha(g)$  and  $\alpha(h)$  conjugate (simultaneously) in  $M_r(\mathbb{Q}\zeta_q)$  to diagonal matrices

$$\begin{aligned}\alpha'(g) &= \text{diag}(\mu_1, \dots, \mu_r), \\ \alpha'(h) &= \text{diag}(v_1, \dots, v_r) \quad (\mu_i, v_i \in \langle \zeta_q \rangle).\end{aligned}$$

Write

$$\Gamma^{-1}(\tfrac{1}{2}(h + h^{-1})) = \sum_j \lambda_j h^j \in (\widehat{\mathbb{Z}}_p[\langle h \rangle])^*.$$

Then, since  $K_2(\widehat{\mathbb{Z}}_p[\chi])$  injects into  $K_2(\widehat{\mathbb{Z}}_p\zeta_q)$  (see [20]),

$$\begin{aligned}\varphi_\chi(\{g, \Gamma^{-1}(\tfrac{1}{2}(h + h^{-1}))\}) &= \left\{ \alpha(g), \sum_j \lambda_j \alpha(h)^j \right\} \\ &= \prod_{i=1}^r \left\{ \mu_i, \sum_j \lambda_j v_i^j \right\} \in K_2(\widehat{\mathbb{Z}}_p\zeta_q).\end{aligned}$$

Also, taking norm residue symbols commutes with inclusions of cyclotomic fields  $\mathbb{Q}_p \zeta_{p^i}$  for odd  $p$  (this follows, e.g., from the formulas in [18]). So by [17, Lemma 1.4],

$$\lambda_\chi \circ \varphi_\chi(\{g, \Gamma^{-1}(\tfrac{1}{2}(h+h^{-1}))\}) = \prod_{i=1}^r \left( \mu_i, \sum_j \lambda_j v_i^j \right)_v = \prod_{i=1}^r \mu_i^{\tau(v_i)},$$

where  $\tau(v_i) = 1$  if  $v_i = 1$ ,  $\tau(v_i) = 0$  if  $v_i \neq 1$ . But the last expression is just  $\det(g, V_\chi^h)$ , and so (1) is proved.

To show (2), fix  $g \in \pi$  and  $u \in (\hat{\mathbb{Z}}_p[\hat{\mathbb{Z}}_\pi(g)])^*$ , and let  $\bar{u}$  be the unit obtained from  $u$  by replacing  $h \mapsto h^{-1}$ . Then

$$\lambda_\chi \circ \varphi_\chi(\{g, u\}) = \lambda_\chi \circ \varphi_\chi(\{g, \bar{u}\}):$$

$(\zeta_{p^k}, a)_v = (\zeta_{p^k}, \bar{a})_v$  for any  $a \in (\hat{\mathbb{Z}}_p \zeta_{p^k})^*$  by [18]. So by (1),

$$\begin{aligned} \lambda_\chi \circ \varphi_\chi(\{g, u\}) &= \lambda_\chi \circ \varphi_\chi(\{g, u\bar{u}\})^{1/2} = \psi_\chi(g \otimes \Gamma(u\bar{u}))^{1/2} \\ &= \psi_\chi(g \otimes \tfrac{1}{2}(\Gamma(u) + \Gamma(\bar{u}))) = \psi_\chi(g \otimes \Gamma(u)): \end{aligned}$$

the last step since  $\psi_\chi(g \otimes h) = \psi_\chi(g \otimes h^{-1})$  ( $V_\chi^h = V_\chi^{h^{-1}}$ ). ■

We use the term a cluster of characters for a  $p$ -group  $\pi$  to mean any set of irreducible characters which contains exactly one character dividing each irreducible  $\mathbb{Q}\pi$ -module. Thus in the above notation, if  $S$  is a cluster of characters for  $\pi$ , then there is a natural isomorphism

$$\mathbb{Q}\pi \cong \prod_{\chi \in S} M_{r_\chi}(\mathbb{Q}\chi).$$

Furthermore, for each  $\chi \in S$ ,

$$K_2^{\text{top}}(\hat{\mathbb{Z}}_p[\chi]) \cong (\mu_{\mathbb{Q}\chi})_p \cong \varprojlim_n SK_1(\mathbb{Z}\chi, p^n)$$

by [20, Sect. 5]. So Proposition 1.8 in [11] takes the form

$$Cl_1(\mathbb{Z}\pi) \cong \text{Coker}(\varphi_S).$$

Since  $\text{Im}(\psi_S) \subseteq \text{Im}(\lambda_S \circ \varphi_S)$  by Proposition 7.1, this implies:

**THEOREM 7.2.** *Let  $p$  be an odd prime, let  $\pi$  be a  $p$ -group, and let  $S$  be a cluster of characters for  $\pi$ . Then  $\lambda_S^{-1}$  induces a surjection*

$$A_S: \text{Coker}(\psi_S) \twoheadrightarrow Cl_1(\mathbb{Z}\pi).$$

The following conjecture is motivated partly since it holds in the abelian case [17, Theorem 1.8], partly by the similarity between  $H_1(\pi; \hat{\mathbb{Z}}_p \pi)$  and

$K_2^*(\hat{\mathbb{Z}}_p\pi)$  (Theorem 3.6), and partly since it seems to be the best hope for finding a “reasonable” combinatorial algorithm for computing  $Cl_1(\mathbb{Z}\pi)$ .

**Conjecture 7.3.** For any odd prime  $p$ , any  $p$ -group  $\pi$ , and any cluster  $S$  of characters for  $\pi$ ,

$$A_S: \text{Coker}(\psi_S) \rightarrow Cl_1(\mathbb{Z}\pi)$$

is an isomorphism.

Note that if Conjecture 7.3 holds, then this together with [11, Theorem 4.8(ii)] and [22, Theorem 2(i)] gives a direct combinatorial algorithm for computing odd torsion in  $SK_1(\mathbb{Z}\pi)$  for an arbitrary finite group  $\pi$ . Hopefully, if Conjecture 7.3 could be shown, then a similar but more complicated algorithm could be found for computing  $Cl_1(\mathbb{Z}\pi)$  when  $\pi$  is a 2-group (see Theorem 1.8 in [17] for the abelian case).

The best we can do so far towards proving Conjecture 7.3 is to show that  $\text{Im}(\lambda_{S'} \circ \varphi_{S'}) = \text{Im}(\psi_{S'})$  when  $S'$  is a set of *linear* characters for  $\pi$ : for example, a cluster of characters for  $\pi^{ab}$ . This does at least give a lower bound for  $Cl_1(\mathbb{Z}\pi)$ , i.e., it is at least as large as  $\text{Coker}(\psi_{S'})$  when  $S'$  is such a cluster of characters.

In fact, this result can be extended to include 2-groups. If  $\pi$  is a 2-group and  $T$  any set of linear characters for  $\pi$ , then

$$\psi_T = \prod \psi_\chi: H_1(\pi; \hat{\mathbb{Z}}_2\pi) \rightarrow \prod_{\chi \in T} (\mu_{\mathbb{Q}\chi})_2$$

is defined by setting, for  $\chi \in T$  and commuting  $g, h \in \pi$ ,

$$\begin{aligned} \psi_\chi(g \otimes h) &= \chi(g) & \text{if } \chi(h) &= 1, \\ &= 1 & \text{if } \chi(h) &\neq 1. \end{aligned}$$

Note that this coincides with the definition of  $\psi_T$  for odd  $p$ -groups—in the case of linear characters.

When  $T$  is a set of linear characters for  $\pi$  such that for all  $\chi \in T$ ,  $\mathbb{Q}\chi = \mathbb{Q}\zeta_{2^n}$  for some  $n \geq 2$ , then a modified norm residue symbol map

$$\lambda'_T = \prod_{\chi \in T} \lambda'_\chi: \prod_{\chi \in T} K_2^{\text{top}}(\hat{\mathbb{Z}}_2[\chi]) \xrightarrow{\cong} \prod_{\chi \in T} (\mu_{\mathbb{Q}\chi})_2 = \prod_{\chi \in T} \text{Im}(\chi)$$

is defined as follows: if  $|\text{Im}(\chi)| = 2^n$  and  $a, b \in (\hat{\mathbb{Z}}_2[\chi])^*$ , then

$$\lambda'_\chi(\{a, b\}) = (a, b)_v^{2^{n-1}+1}.$$

To simplify the statement of the next theorem, we set  $\lambda'_T = \lambda_T$  whenever  $T$  is a set of characters for an odd  $p$ -group.

Finally, for any 2-group  $\pi$ , let

$$v: H_1(\pi; \hat{\mathbb{Z}}_2\pi) \rightarrow H_2(\pi)$$

be the map defined in Section 3:  $v(g \otimes h) = g \wedge h \in H_2(\pi)$  for any pair of commuting elements  $g, h \in \pi$ .

**THEOREM 7.4.** *Let  $p$  be any prime, let  $\pi$  be a  $p$ -group, and let  $T$  be a cluster of characters for  $\pi^{ab}$ . Set  $T_0 = T$  if  $p$  is odd;*

$$T_0 = \{\chi \in T: |\text{Im}(\chi)| > 2\} = \{\chi \in T: \mathbb{Q}\chi \not\subseteq \mathbb{R}\}$$

*if  $p = 2$ . Consider the maps*

$$\begin{aligned} K_2^{\text{top}}(\hat{\mathbb{Z}}_p\pi) &\xrightarrow{\lambda'_{T_0} \circ \varphi_{T_0}} \prod_{\chi \in T_0} (\mu_{\mathbb{Q}\chi})_p = \prod_{\chi \in T_0} \text{Im}(\chi) \\ &\xleftarrow{\psi_{T_0}} H_1(\pi; \hat{\mathbb{Z}}_p\pi). \end{aligned}$$

(i) *If  $p$  is odd, then  $\text{Im}(\lambda'_T \circ \varphi_T) = \text{Im}(\psi_T)$ , and there is a surjection*

$$Cl_1(\mathbb{Z}\pi) \twoheadrightarrow \left[ \prod_{\chi \in T} \text{Im}(\chi) \right] / \psi_T(H_1(\pi; \hat{\mathbb{Z}}_p\pi)).$$

(ii) *If  $p = 2$ , then*

$$\text{Im}(\lambda'_{T_0} \circ \varphi_{T_0}) = \psi_{T_0}(\text{Ker}[v: H_1(\pi; \hat{\mathbb{Z}}_2\pi) \rightarrow H_2(\pi)]),$$

*and there is a surjection*

$$Cl_1(\mathbb{Z}\pi) \twoheadrightarrow \left[ \prod_{\chi \in T_0} \text{Im}(\chi) \right] / \psi_{T_0}(\text{Ker}[v: H_1(\pi; \hat{\mathbb{Z}}_2\pi) \rightarrow H_2(\pi)]).$$

*Proof.* Set

$$X = \langle \psi_{T_0}(g \otimes g^m): g \in \pi, m \in \mathbb{Z} \rangle \subseteq \prod_{\chi \in T_0} \text{Im}(\chi).$$

We first claim that

$$X \subseteq \psi_{T_0}(\text{Ker}(v)) \cap \text{Im}(\lambda'_{T_0} \circ \varphi_{T_0}) \quad (1)$$

and that

$$\langle \lambda'_{T_0} \circ \varphi_{T_0}(\{u, v\}): u, v \in (\hat{\mathbb{Z}}_p\sigma)^*, \text{ some cyclic } \sigma \subseteq \pi \rangle \subseteq X. \quad (2)$$

By the naturality of  $\psi$  and  $\lambda' \circ \varphi$  (see the remarks before Proposition 1.5 in

[17]), it suffices to prove (1) and (2) for cyclic  $\pi$ . But in this case,  $H_2(\pi) = 0$ ;  $\psi_{T_0}$  and  $\varphi_{T_0}$  are onto since  $SK_1(\mathbb{Z}\pi) = 0$  (use Theorem 1.8 in [17]);  $X = \text{Im}(\psi_{T_0})$  by definition; and the result follows.

Now consider the following diagram:

$$\begin{array}{ccccc} K_2^{\text{top}}(\hat{\mathbb{Z}}_p \pi) & \xrightarrow{K_2(\alpha)} & K_2^{\text{top}}(\hat{\mathbb{Z}}_p[\pi^{ab}]) & \xrightarrow{\bar{\varphi}} & \left[ \prod_{\chi \in T_0} \text{Im}(\chi) \right] / X \\ \downarrow \Gamma_2^* & & \downarrow \Gamma_2^* & & \\ \text{Ker}(v) \rightarrow \mathcal{U}_0(\pi) & \xrightarrow{\mathcal{U}_0(\alpha)} & \mathcal{U}_0(\pi^{ab}) & \xrightarrow{\bar{\psi}} & \end{array} \quad (3)$$

Here,  $\bar{\varphi}$  and  $\bar{\psi}$  are the reductions of the maps  $\lambda' \circ \varphi$  and  $\psi$  for  $\pi^{ab}$ ; recall that

$$\mathcal{U}(\pi^{ab}) = H_1(\pi^{ab}; \hat{\mathbb{Z}}_p[\pi^{ab}]) / \langle g \otimes g^m : g \in \pi^{ab}, m \in \mathbb{Z} \rangle.$$

Also,  $\alpha: \pi \rightarrow \pi^{ab}$  denotes the projection, and  $K_2(\alpha)$  and  $\mathcal{U}_0(\alpha)$  are the induced maps. The square in (3) commutes by naturality of  $\Gamma_2^*$ .

By Corollary 1.2 in [17],  $K_2^{\text{top}}(\hat{\mathbb{Z}}_p[\pi^{ab}])$  is generated by symbols  $\{1 + p, 1 + px\}$  ( $x \in \hat{\mathbb{Z}}_p[\pi^{ab}]$ ),  $\{g, \lambda\}$  ( $g \in \pi^{ab}$ ,  $\lambda \in (\hat{\mathbb{Z}}_p)^*$ ), and  $\{g, u\}$  ( $g \in \pi^{ab}$ ,  $u \in 1 + I(\hat{\mathbb{Z}}_p[\pi^{ab}])$ ). That

$$\{1 + p, 1 + px\}, \{g, \lambda\} \in \text{Ker}(\bar{\varphi}) \cap \text{Ker}(\Gamma_2^*)$$

follows from (2) and the fact that  $\Gamma_2^*$  factors through  $K_2^*(\hat{\mathbb{Z}}_p[\pi^{ab}])$  ( $K_1(\hat{\mathbb{Z}}_p \pi, p)$  is generated by cyclic induction by, e.g., [10, Proposition 2]). On the other hand, for any  $g$  and  $u$ ,

$$\Gamma_2^*(\{g, u\}) = g \otimes \Gamma(u)$$

by Proposition 3.8, and by Lemma 1.4 in [17],

$$\bar{\varphi}(\{g, u\}) = \bar{\psi}(g \otimes \Gamma(u)).$$

So the triangle in diagram (3) commutes.

Hence,

$$\begin{aligned} \text{Im}(\lambda'_{T_0} \circ \varphi_{T_0}) / X &= \text{Im}(\bar{\varphi} \circ K_2(\alpha)) = \text{Im}(\bar{\psi} \circ \mathcal{U}_0(\alpha)) \\ &= \psi_{T_0}(\text{Ker}(v)) / X. \end{aligned}$$

Together with (1), this shows that

$$\text{Im}(\lambda'_{T_0} \circ \varphi_{T_0}) = \psi_{T_0}(\text{Ker}(v)). \quad (4)$$

So we are done if  $p = 2$ . If  $p$  is odd, then  $\text{Im}(\lambda_{T'} \circ \varphi_T) \supseteq \text{Im}(\psi_T)$  by Proposition 7.1, and so they are equal by (4). ■

Theorem 7.4 provides a means of detecting elements in  $Cl_1(\mathbb{Z}\pi)$  which vanish in  $Cl_1(\mathbb{Z}[\pi^{ab}])$ . These are precisely the elements which were detected in Propositions 16 and 18 in [21], and in Step 7 of the proof of Theorem 5.6 in [11]. In fact, Theorem 7.4 and its proof give some explanation of why the proofs in Section 2 of [21] follow such a regular pattern.

We end with the following example of the use of Theorems 7.2 and 7.4:

**THEOREM 7.5.** *For any odd prime  $p$  and any non-abelian group  $\pi$  of order  $p^3$ ,*

$$SK_1(\mathbb{Z}\pi) = Cl_1(\mathbb{Z}\pi) \cong (\mathbb{Z}/p)^{p-1}.$$

*Proof.* That  $SK_1(\hat{\mathbb{Z}}_p\pi) = 0$  (and hence  $SK_1(\mathbb{Z}\pi) = Cl_1(\mathbb{Z}\pi)$ ) follows from Proposition 9 in [10].

Note that  $\pi^{ab} \cong \mathbb{Z}/p \times \mathbb{Z}/p$ . Fix  $a, b \in \pi$  which generate  $\pi^{ab}$ , then  $z = [a, b]$  generates  $[\pi, \pi]$ . Furthermore,

$$\mathbb{Q}\pi \cong \mathbb{Q}[\pi^{ab}] \times M_p(\mathbb{Q}\zeta_p) \cong \mathbb{Q} \times (\mathbb{Q}\zeta_p)^{p+1} \times M_p(\mathbb{Q}\zeta_p).$$

So there is a cluster  $S$  of characters for  $\pi$  of the form

$$S = \{1, \chi_0, \dots, \chi_p, \bar{\chi}\},$$

where  $\bar{\chi}$  is a non-linear character and  $\text{Im}(\chi_i) = \langle \zeta_p^i \rangle$  for  $0 \leq i \leq p$ . Set  $T = S - \{\bar{\chi}\}$ , a cluster of characters for  $\pi^{ab}$ .

By Theorems 7.2 and 7.4, there are surjections

$$\text{Coker}(\psi_S) \twoheadrightarrow Cl_1(\mathbb{Z}\pi) \twoheadrightarrow \text{Coker}(\psi_T).$$

On the other hand,  $\psi_\chi(z \otimes a) = 1$  for any  $\chi \in T$  (since  $\chi(z) = 1$ ),  $z$  acts on  $V_{\bar{\chi}}$  via multiplication by  $\zeta$  for some primitive  $p$ th root of unity  $\zeta$ , and so  $\psi_{\bar{\chi}}(z \otimes a) = \zeta$  since  $V_{\bar{\chi}}^a$  is one-dimensional. It follows that

$$\text{Coker}(\psi_S) \cong \text{Coker}(\psi_T) \cong Cl_1(\mathbb{Z}\pi).$$

Finally,  $\psi_T$  factors as a composite

$$\psi_T: H_1(\pi; \hat{\mathbb{Z}}_p\pi) \rightarrow H_1(\pi^{ab}; \hat{\mathbb{Z}}_p[\pi^{ab}]) \rightarrow \prod_{\chi \in T} \text{Im}(\chi) \cong (\mathbb{Z}/p)^{p+1}.$$

Since  $H_1(\pi; \hat{\mathbb{Z}}_p\pi)$  is generated by elements  $g \otimes h$  for commuting  $g, h \in \pi$ , it follows that  $\text{Im}(\psi_T)$  is generated by  $\psi_T(a \otimes e)$  and  $\psi_T(b \otimes e)$ . These are clearly independent, and so

$$Cl_1(\mathbb{Z}\pi) \cong \text{Coker}(\psi_T) \cong (\mathbb{Z}/p)^{p-1}. \quad \blacksquare$$

*Note added in proof.* (1) Conjecture 7.3—the proposed algorithm for describing  $Cl_1(\mathbb{Z}\pi)$  when  $\pi$  is a  $p$ -group and  $p$  is odd—has recently been proven.

(2) When  $p$  is any prime,  $\pi$  is an abelian  $p$ -group, and  $A$  is an unramified  $p$ -ring, a short exact sequence

$$0 \rightarrow K_2^{\text{lop}}(A\pi)/\langle \{\pm g, \pm h\} \rangle \xrightarrow{\Gamma_2} \frac{\pi \otimes A\pi}{\langle g \otimes \lambda g \rangle} \xrightarrow{\hat{v}} \frac{\pi \otimes \pi}{\langle g \otimes h + h \otimes g \rangle} \rightarrow 0$$

has now been constructed. Here,  $\Gamma_2$  and  $\hat{v}$  are liftings of the homomorphisms  $\Gamma_2^*$  and  $v$  constructed in Section 3.

## REFERENCES

1. A. BAK AND U. REHMANN, The congruence subgroup and metaplectic problems for  $SL_{n \geq 2}$  of division algebras, *J. Algebra* **78** (1982), 475–547.
2. H. BASS, “Algebraic  $K$ -Theory,” Benjamin, New York, 1968.
3. H. CARTAN AND S. EILENBERG, “Homological Algebra,” Princeton Univ. Press, Princeton, N. J., 1956.
4. R. K. DENNIS, M. KEATING, AND M. STEIN, Lower bounds for the order of  $K_2(\mathbb{Z}G)$  and  $Wh_2(G)$ , *Math. Ann.* **223** (1976), 97–103.
5. D. GORENSTEIN, “Finite Groups,” Harper & Row, New York, 1968.
6. P. HILTON AND U. STAMMBACH, “A course in Homological Algebra,” Springer-Verlag, New York/Berlin, 1971.
7. M. KEATING, Values of tame symbols on division algebras, *J. London Math. Soc.* **14** (1976), 25–30.
8. J.-L. LODAY,  $K$ -Théorie algébrique et représentations de groupes, *Ann. Sci. Ecole Norm. Sup.* **9** (1976), 309–377.
9. J. MILNOR, “Introduction to Algebraic  $K$ -Theory,” Princeton Univ. Press, Princeton, N. J., 1971.
10. R. OLIVER,  $SK_1$  for finite group rings, II, *Math. Scand.* **47** (1980), 195–231.
11. R. OLIVER,  $SK_1$  for finite group rings, IV, *Proc. London Math. Soc.* **46** (1983), 1–37.
12. U. STAMMBACH, “Homology in Group Theory,” Lecture Notes in Mathematics No. 359, Springer-Verlag, New York/Berlin, 1973.
13. M. STEIN, Surjectivity stability in dimension 0 for  $K_2$  and related functors, *Trans. Amer. Math. Soc.* **178** (1973), 165–191.
14. C. T. C. WALL, On the classification of hermetian forms, III. Complete semilocal rings, *Invent. Math.* **19** (1973), 59–71.
15. A. WEIL, “Basic Number Theory,” Springer-Verlag New York/Berlin, 1967.
16. E. WEISS, “Cohomology of Groups,” Academic Press, New York, 1969.
17. R. ALPERIN, R. K. DENNIS, R. OLIVER, AND M. STEIN,  $SK_1$  of finite abelian groups, II, to appear.
18. E. ARTIN AND H. HASSE, Die beiden Ergänzungssätze zum Reziprozitätsgesetz der  $l^n$ -ten Potenzreste im Körper der  $l^n$ -ten Einheitswurzeln, *Abh. Math. Sem. Univ. Hamburg* **6** (1928), 146–162.
19. S. BLOCH, Algebraic  $K$ -theory and crystalline cohomology, *I.H.E.S. Publ. Math.* **47** (1977), 187–268.
20. R. K. DENNIS AND M. STEIN,  $K_2$  of discrete valuation rings, *Adv. in Math.* **18** (1975), 182–238.
21. R. OLIVER,  $SK_1$  for finite group rings, I, *Invent. Math.* **57** (1980), 183–204.
22. R. OLIVER,  $SK_1$  for finite group rings, III, in “Lecture Notes in Mathematics No. 854,” pp. 299–337, Springer-Verlag, New York/Berlin, 1981.
23. D. QUILLLEN, Higher algebraic  $K$ -theory, I, in “Lecture Notes in Mathematics No. 341,” pp. 85–147, Springer-Verlag, New York/Berlin, 1973.
24. P. ROQUETTE, Realisierung von Darstellungen endlicher nilpotenter Gruppen, *Ark. Math.* **9** (1958), 241–250.